

GENOTYPING BY PCR PROTOCOL
MUTANT MOUSE RESOURCE & RESEARCH CENTER: UC DAVIS

mmrrc@ucdavis.edu

530-754-MMRRC

UCD_050581Geno_ProtocolProtocol

FVB.129P2-Kat5tm1Tfaz/Mmucd

MMRRC: 043817-

Protocol:

Reagent/Constituent	Volume (μ L)
Water	5.6
GoTaq® G2 Colorless Master Mix, 2X	7.5
Primer 1. (stock concentration is 20 μ M) comF	0.45
Primer 2. (stock concentration is 20 μ M) wtR	0.45
DNA (example) extracted w/ "Qiagen DNeasy columns or other similar silica based kits"	1.0
	TOTAL VOLUME

Comments on protocol:

- Protocol may work with other DNA extraction methods.
- Use Touch-Down cycling protocol-first 10 cycles anneal at 65°C decreasing in temperature by 1.0°C; next 30 cycles anneal at 55°C.

Strategy:

Steps	HOT START? <input type="checkbox"/>	Temp (°C)	Time (m:ss)	# of Cycles
1. Initiation/Melting		94	5:00	1x
2. Denaturation		94	0:15	
3. Annealing	steps 2-3-4 cycle in sequence	65 ($\downarrow 1^{\circ}\text{C}/\text{cycle}$)	0:30	10x
4. Elongation		72	0:40	
5. Denaturation		94	0:15	
6. Annealing	steps 5-6-7 cycle in sequence	55	0:30	30X
7. Elongation		72	0:40	

Primers:

Electrophoresis Protocol:

Name	Nucleotide Sequence (5' - 3')	Agarose: 1.5%	V: 90
1. 43817-F	GTGGGCTACTTCTCCAAGGTC	Estimated Running Time: 90 min.	
2. 43817-R	GCAGCGCATGCCCTCTATC	Primer Combination	Band (bp)
		1 & 2	320
		1 & 2	280
			wildtype
			mutant

