

GENOTYPING BY PCR PROTOCOL

MUTANT MOUSE RESOURCE & RESEARCH CENTER: UC DAVIS

mmrrc@ucdavis.edu

530-754-MMRRC

Protocol Name: C57BL/6NTac-Brd1tm1.2Arte/AborgMmucd **MMRRC:** 065563-UCD

Protocol:

Reagent/Constituent	Volume (μL)
Water	4.7
GoTaq® G2 Colorless Master Mix, 2X	7.5
Primer 1. (stock concentration is 20μM)	0.45
Primer 2. (stock concentration is 20μM)	0.45
Primer 3. (stock concentration is 20μM)	0.45
Primer 4. (stock concentration is 20μM)	0.45
DNA (example) extracted w/ "Qiagen DNeasy columns or other similar silica based kits"	1.0
TOTAL VOLUME	
15	

Comments on protocol:

- Protocol may work with other DNA extraction methods.
- Use Touch-Down cycling protocol-first 10 cycles anneal at 65°C decreasing in temperature by 1.0°C; next 30 cycles anneal at 55°C.

Strategy:

Steps	Temp (°C)	Time (m:ss)	# of Cycles
1. Initiation/Melting HOT START? ☐	94	2:00	1x
2. Denaturation	94	0:10	
3. Annealing steps 2-3-4 cycle in sequence	65 (↓ 1°C/cycle)	0:30	10x
4. Elongation	68	2:00	
5. Denaturation	94	0:15	
6. Annealing steps 5-6-7 cycle in sequence	55	0:30	30x
7. Elongation	68 (+20s/cycle)	2:00	

Primers:

Electrophoresis Protocol:

Name	Nucleotide Sequence (5' - 3')	Argarose: 1.5% V: 90		
1. 65563-mutF	ATAAACACCTGCAAGCATCC	Estimated Running Time: 90 min.		
2. 65563-mutR	GTAAGAGTACCGTGGTTAGC	Primer Combination	Band (bp)	Genotype
3. 65563-conF	GTGGCACGGAACCTTCTAGTC	1 & 2	443	mutant
4. 65563-conR	CTTGTCAGTAGCAGGAAGA	3 & 4	335	Control pcr
		1 & 2	2344	wildtype

