

GENOTYPING PROTOCOL

MUTANT MOUSE RESOURCE & RESEARCH CENTER: UC DAVIS

mmrrc@ucdavis.edu

530-754-MMRRC

Protocol Name: CR10522 Cln8 iDex Stock # 065590

Protocol: GoTaq® G2 Colorless Master Mix(Promega)

Reagent/Constituent	Volume (μL)
Water	4.5
GoTaq® G2 Colorless Master Mix,2X	7.5
Primer 1. (stock concentration is 20μM) comF	0.5
Primer 2. (stock concentration is 20μM) wtR	0.5
Primer 3. (stock concentration is 20μM) mutR	0.5
DNA (example) extracted w/ "Qiagen DNeasy columns or other similar silica based kits"	1.5
TOTAL VOLUME OF REACTION:	15.00 μL

Comments on protocol:

- Protocol may work with other DNA extraction methods.

Strategy:

Steps	Temp (°C)	Time (m:ss)	# of Cycles
1. Initiation/Melting HOT START? ☐	94	2:00	1x
2. Denaturation	94	0:10	
3. Annealing steps 2-3-4 cycle in sequence	65 (↓1°C/cycle)	0:30	10x
4. Elongation	68	2:00	
5. Denaturation	94	0:15	
6. Annealing steps 5-6-7 cycle in sequence	55	0:30	25x
7. Elongation	68	2:00 (↑20sec/cycle)	
8. Finish	4	∞	n/a

Primers:

Name	Nucleotide Sequence (5' - 3')	Agarose: 1.5% V: 90
1. CR_Cln8_XD_A67V_comF	AGATTCAGATCATGACTCCTGTGAGCA	Estimated Running Time: 90 min.
2. CR_Cln8_XD_A67V_wtR*	CGAAGACAGCACGCGTCGC	Primer Combination Band (bp) Genotype
3. CR_Cln8_XD_A67V_mutR	GGTGGTGGACAACCAGAAACAAGTC	1 & 2, 1 & 3 231,429 wildtype
		1 & 3 376 mutant

Electrophoresis Protocol:

Allele Description: Exon 2 [ENSMUSE00000270532](#) had 53bp deleted from the 183rd coding nucleotide through the 238th coding nucleotide from the Cln8 gene [ENSMUSG0000026317.7](#) using CRISPR Cas9 gene editing technology in mouse zygotes. Subsequent founders were backcrossed to C57BL6/N to produce sequence confirmed heterozygous animals.

*wtR primer untested (ePCR verified)

