

GENOTYPING BY PCR PROTOCOL

MUTANT MOUSE RESOURCE & RESEARCH CENTER: UC DAVIS

mmrrc@ucdavis.edu

530-754-MMRRC

Protocol Name: B6J.B6N(Cg)-Aldh1a1tm1.1(Aldh1a1/cre/ERT2)Cai/Mmucd **MMRRC:** 065802-UCD

Protocol:

Reagent/Constituent	Volume (μL)
Water	5.6
GoTaq® G2 Colorless Master Mix, 2X	7.5
Primer 1. (stock concentration is 20μM)	0.45
Primer 2. (stock concentration is 20μM)	0.45
DNA (example) extracted w/ "Qiagen DNeasy columns or other similar silica based kits"	1.0
TOTAL VOLUME	
15	

Comments on protocol:

- Protocol may work with other DNA extraction methods.
- Use Touch-Down cycling protocol-first 10 cycles anneal at 65°C decreasing in temperature by 1.0°C; next 30 cycles anneal at 55°C.

Strategy:

Steps	Temp (°C)	Time (m:ss)	# of Cycles
1. Initiation/Melting HOT START? ☐	94	5:00	1x
2. Denaturation	94	0:15	
3. Annealing steps 2-3-4 cycle in sequence	65 (↓1°C/cycle)	0:30	10x
4. Elongation	72	0:40	
5. Denaturation	94	0:15	
6. Annealing steps 5-6-7 cycle in sequence	55	0:30	30X
7. Elongation	72	0:40	

Primers:

Electrophoresis Protocol:

Name	Nucleotide Sequence (5' - 3')	Argarose: 1.5% V: 90		
1. 65802-CreF	GCGCGCCTGAAGATATAGAAGA	Estimated Running Time: 90 min.		
2. 65802-wtF	CTTCAAGGAGTCACTGAGGGAC	Primer Combination	Band (bp)	Genotype
3. 65802-comR	GCTCAAGACAGTCGCAATGAAG	1 & 3	~320	mutant
		2 & 3	394	wildtype

