GENOTYPING PROTOCOL

MUTANT MOUSE RESOURCE & RESEARCH CENTER: UC DAVIS

mmrrc@ucdavis.edu
530-754-MMRRC

Protocol Name: **CR10387 Arhgef26 EXDEL em2(IMPC)MBP**

Protocol: ***GoTaq® G2 Colorless Master Mix(Promega)***

Reagent/Constituent	Volume (µL)
Water	4.5
GoTaq® G2 Colorless Master Mix,2X	7.5
Primer 1. (stock concentration is 20µM) comF	0.5
Primer 2. (stock concentration is 20µM) wtR	0.5
Primer 3. (stock concentration is 20µM) mutR	0.5
DNA (example) extracted w/ "Qiagen DNeasy columns or other similar silica based kits"	1.5
TOTAL VOLUME OF REACTION:	**15.00 µL**

Comments on protocol:

- Protocol may work with other DNA extraction methods.

Strategy:

Steps		Temp (ºC)	Time (m:ss)	# of Cycles
1. Initiation/Melting	HOT START? ☐	94	2:00	**1x**
2. Denaturation		94	0:10	
3. Annealing	steps 2-3-4 cycle in sequence	65 (↓1ºC/cycle)	0:30	**10x**
4. Elongation		68	2:00	
5. Denaturation		94	0:15	
6. Annealing	steps 5-6-7 cycle in sequence	55	0:30	**25x**
7. Elongation		68	2:00 (↑20sec/cycle)	
8. Finish		4	∞	n/a

Primers:

Name	Nucleotide Sequence (5' - 3')
1. CR_Arhgef26_comF	GCTTCATGGACATTGAGACTTCTCG
2. CR_Arhgef26_wtR*	CCTCAATGTCTTTGTCAAGTACCCATG
3. CR_Arhgef26_mutR	GCATTCTATAAGGCTGGAAGGCTGA

Electrophoresis Protocol:

Agarose: **1.5%** **V:** **90**

Estimated Running Time: **90** min.

Primer Combination	**Band** (bp)	**Genotype**
1 & 2,1 & 3	341,1375	wildtype
1 & 3	623	mutant

Allele Description: Exon 3 ENSMUSE00001222548 and flanking splicing regions were constitutively deleted from the Arhgef26 gene ENSMUSG00000036885.14 using CRISPR Cas9 gene editing technology in mouse zygotes. Subsequent founders were backcrossed to C57BL6/N to produce sequence confirmed heterozygous animals.

*wtR primer untested (ePCR verified)