

GENOTYPING BY PCR PROTOCOL

MUTANT MOUSE RESOURCE & RESEARCH CENTER: UC DAVIS

mmrrc@ucdavis.edu

530-754-MMRRC

Protocol Name: B6J.Cg-Tg(Eif2s3x)6Mjmm/Mmucd MMRRC: 043568-UCD

Protocol:

Reagent/Constituent	Volume (µL)
Water	5.6
GoTaq® G2 Colorless Master Mix, 2X	7.5
Primer 1. (stock concentration is 20µM)	0.45
Primer 2. (stock concentration is 20µM)	0.45
DNA (example) extracted w/ "Qiagen DNeasy columns or other similar silica based kits"	1.0
TOTAL VOLUME	
15	

Comments on protocol:

Strategy:

Steps	Temp (°C)	Time (m:ss)	# of Cycles
1. Initiation/Melting HOT START? <input type="checkbox"/>	94	5:00	1x
2. Denaturation	94	0:15	
3. Annealing steps 2-3-4 cycle in sequence	65 (↓1°C/cycle)	0:30	10x
4. Elongation	72	0:40	
5. Denaturation	94	0:15	
6. Annealing steps 5-6-7 cycle in sequence	55	0:30	30X
7. Elongation	72	0:40	

Primers:

Electrophoresis Protocol:

Name	Nucleotide Sequence (5' - 3')	Argarose: 1.5%	V: 90
1. 43568-pTARBAC1	CGCAGCCGTGTAACCGAGCATAG	Estimated Running Time: 90 min.	
2. 43568-pTARBAC2	AAAAAGAAAGGCCGCCAGACGACT	Primer Combination	Band (bp)
3. 43568-PTARBAC-F	GCCTGATGCGGTATTTTCTC	1 & 2	480
4. 43568-PTARBAC-R	CGGCATCAGAGCAGATTGTA	3 & 4	87
			Genotype
			mutant
			mutant

