

Culturing MMRRC-Distributed Lex1 Clones

Cell Line Information

The MMRRC-distributed Lexicon derived (Lex1) clones are feeder-dependent.

Parental ES cells: Lex1, isolated from 129SvEvBrd mouse strain
Injection of cells into C57BL/6 Albino blastocysts will produce agouti/white chimeras.

Feeder cells: MEF feeder cells express very small amounts of Leukemia Inhibitory Factor (LIF) and neomycin phosphotransferase (*Neo*). This cell line is mycoplasma free.

Reagents and Supplies

<u>Item</u>	<u>Vendor</u>	<u>Catalog Number</u>
DMEM, high glucose, with glutamine	Gibco	11965-092
Penicillin/Streptomycin, 100 u/ml	Gibco	15140-122
L-Glutamine	Gibco	25030-081
Trypsin EDTA (1X, 0.25%) (supplement the 0.25% with 1% Chick Serum)	Gibco	25300-054 (0.05%) 25200-072 (0.25%)
LIF (ESGRO)	Gibco	13275-029
Fetal Bovine Serum, Defined	Hyclone	SH30070.03
PBS (1X without Ca or Mg)	Gibco	14190-144
DMSO, 100 ml (2X Freezing medium: Lexicon Cell Medium with 20% DMSO - make fresh as required)	Sigma	D2650
Mitocycin C 10x2mg (10µg/ml Inactivation media: add 2mg of Mitomycin C to 200 ml MEF Feeder medium -may be stored at -20°C for up to 6 months)	Sigma	M0503
2(β)-Mercaptoethanol (1000x working soln: add 70µl 2-Mercaptoethanol to 9.93ml PBS. Store at 4°C, and make fresh every 2 weeks)	Sigma	M-7522
Gelatin, 2% (0.1% working soln: add 25 ml of 2% solution to 475ml of PBS. Store at 4°C)	Sigma	G1393

MEF Feeder Medium (sterile filter through 0.2µM filter unit)

<u>Reagent</u>	<u>Stock Conc.</u>	<u>Final conc.</u>	<u>Total: 500ml</u>
DMEM		1x	435ml
FBS	100%	10%	50ml
L-Glutamine	200mM	2mM	5ml
Sodium Pyruvate	100mM	1mM	5ml
Pen/Strep	10,000U/ml	100U/ml	5ml

MMRRC AB1 Medium (sterile filter through 0.2µM filter unit)

<u>Reagent</u>	<u>Stock Conc.</u>	<u>Final conc.</u>	<u>Total: 597.5ml</u>
DMEM		1x	500ml
FBS	100%	15%	90ml
L-Glutamine	200mM	2mM	3ml
Pen/Strep	10,000U/ml	5,000U/ml	3ml
1000x βME	5.5×10^{-6} M	1µM	0.5ml
LIF	10^7 U/ml	1000U/ml	1ml

MEF Inactivation Media (sterile filter through 0.2µM filter unit)

<u>Reagent</u>	<u>Stock Conc.</u>	<u>Final Conc.</u>	<u>Volume</u>
MEF Feeder medium	1x	1x	200ml
Mitomycin C	2mg powder	10ug/ml	2mg
Total Volume			200 ml

Preparing MEF Feeder Cells

Thawing, Expanding and Treating Active MEF Cells

1. Thaw 1 vial of MEF cells (approximately $1.5-2 \times 10^6$ cells/vial) in a 37°C water bath and dilute into 10 ml of pre-warmed MEF feeder medium.
2. Pellet the cells by spinning for 4 minutes at 1000 rpm in a bench-top clinical centrifuge.
3. Aspirate off medium and gently resuspend cells in 5 ml of pre-warmed MEF feeder medium.
4. Transfer cell suspension to a 6 cm gelatinized dish, and grow at 37°C in a humidified 5% CO₂ incubator (should be confluent within 2-3 days).
5. When confluent, aspirate medium off and wash with 5 ml of pre-warmed PBS, pipetting it away from the cells. Rock dish gently and aspirate medium. Repeat.
6. Cover cells with 1 ml of 0.05% trypsin solution and incubate at 37°C for 4 minutes or until cells are uniformly dispersed into small clumps.
7. Add 5 ml of MEF Feeder medium to inactivate the trypsin, and pipette vigorously to make single cell suspension (we recommend 15 times).
8. Spin for 4 minutes at 1000 rpm.

9. Aspirate off medium and gently resuspend cells in 20 ml of pre-warmed MEF Feeder medium.
10. Split the cell suspension onto two gelatinized 10 cm tissue culture dishes, and grow at 37°C in a humidified 5% CO₂ incubator (should be confluent within 2-3 days).
11. To mitotically inactivate, replace medium with 10 ml Inactivation medium (2mg of Mitomycin C to 200 ml MEF feeder medium), and incubate in a 37°C humidified 5% CO₂ incubator for 2.5 hours. Aspirate Inactivation medium, and rinse three times with pre-warmed PBS; aspirating completely between rinses. These dishes are now ready to use.
12. If you wish to freeze the cells for later usage, trypsinize and pellet the cells as before, but with 1.5 ml of 0.05% trypsin solution, and inactivate the trypsin with 5 ml medium.
13. For each 10 cm dish, count cells, and resuspend in an equal volume of MEF Feeder medium and 2X Freezing medium; to a density of 1.5-2 x 10⁶ cells/0.5 ml. Decant 0.5 ml aliquots into labeled cryovials.
14. Immediately place cryovials in a Styrofoam container or temperature controlled freezing vessel.
15. Freeze vials in a -80°C freezer. After 24 hours, transfer cryovials to liquid or vapor-phase nitrogen for longer term storage.

Plating Mitotically Inactive MEF Feeder Cells

1. Coat a 6 cm tissue culture dish with 0.1% gelatin and aspirate off immediately before use.
2. Thaw 1 vial of mitotically inactive MEF Feeder cells (approx. 1.5-2 x 10⁶ cells) in a 37°C water bath and dilute into 10 ml of pre-warmed MEF feeder cell medium.
3. Pellet the cells by spinning for 4 minutes at 1000 rpm.
4. Aspirate off medium and gently resuspend cells in 5 ml of pre-warmed MEF Feeder cell medium.
5. Transfer cell suspension to the 6 cm gelatinized dish, and grow at 37°C in a humidified 5% CO₂ incubator.
6. The feeders are ready for use after 6-12 hours, or may be maintained in the incubator for a maximum of 8-10 days.

Thawing MMRRC-distributed AB1 clones

1. Thaw 1 vial of ES cells (approximately 5 x 10⁶ cells/vial) in a 37°C water bath and dilute (drop wise) into 10 ml of pre-warmed MMRRC AB1 medium.
2. Pellet the cells by spinning for 4 minutes at 1000 rpm.
3. Aspirate off medium and gently resuspend cells in 5 ml of pre-warmed MMRRC AB1 medium.

4. Aspirate the old medium from your 6 cm mitotically inactive MEF Feeder dish.
5. Transfer the ES cell suspension to the feeder dish, and grow in a 37°C humidified 5% CO₂ incubator.
6. Change medium the following day to remove dead cells and residual DMSO.
7. Change medium daily until 80% confluent (approx. $1.5-2 \times 10^7$ cells); should take 2-3 days.
8. When confluent, the 6 cm dish may be split in two; half for microinjection and half to expand for freezing.

Expansion of MMRRC-distributed AB1 clones for Microinjection and Future Use

1. 1 day prior, prepare one 10 cm MEF Treated Feeder dish, using the plating density guide below.
2. The next day, aspirate off the old medium prior to plating ES cells.
3. On the day, wash the confluent 6 cm ES cell dish once with 5 ml PBS.
4. Cover the cells with 1 ml of 0.25% trypsin solution and incubate at 37°C for 4-5 minutes or until cells are uniformly dispersed into small clumps.
5. Add 5 ml of MMRRC AB1 medium to inactivate the trypsin, and pipette vigorously to make single cell suspension (we recommend 10-15 times).
6. Split the cell suspension in half, placing 2.5 ml each into 15 ml centrifuge tubes (labeled 'Expansion' and 'Microinjection').
7. Spin both tubes for 4 minutes at 1000 rpm.
8. For the 'Expansion' cells; aspirate off the supernatant and resuspend the pellet in 10 ml MMRRC AB1 medium. Transfer the cell suspension onto the 10 cm MEF Treated Feeder dish prepared the day before. Grow in a 37°C humidified 5% CO₂ incubator. Change medium daily until 80% confluent (should take 2-3 days).
9. For the 'Microinjection' cells; aspirate off the supernatant and resuspend the pellet in 150-400 µl microinjection medium (Hepes Buffered DMEM with 5% FBS; filtered through 0.2 µm filter unit, can be aliquoted and stored at -20°C for up to 12 months). Immediately place the cell suspension on ice, and microinject within 1-2.5 hours.

Freezing MMRRC-distributed AB1 clones

1. Wash the confluent 10 cm ES cell dish once with 10 ml PBS each.
2. Cover the cells with 1.5 ml of 0.25% trypsin solution and incubate at 37°C for 4-5 minutes or until cells are uniformly dispersed into small clumps.
3. Add 5 ml MMRRC AB1 medium to inactivate the trypsin, and pipette vigorously to make single cell suspension (we recommend 10-15 times).

4. Spin for 4 minutes at 1000 rpm.
5. Aspirate supernatant and resuspend the pellet in an equal volume of MMRRC AB1 medium and 2X Freezing medium (we would recommend 8-10 vials containing 0.5 ml aliquots; per 10 cm dish). Decant into labeled cryovials.
6. Immediately place cryovials in a Styrofoam container or temperature controlled freezing vessel.
7. Freeze vials in a -80°C freezer. After 24 hours, transfer cryovials to liquid or vapor-phase nitrogen for longer term storage.

MEF Treated Feeder Cells – Plating Density Guide

Plate/Dish	Feeder Cell Density
96 well	1.5×10^4
24 well	1×10^5
3.5 cm dish	8×10^5
6 cm dish	$1.5-2 \times 10^6$
100 mm dish	$3.5-4 \times 10^6$