

GENOTYPING BY PCR PROTOCOL
MUTANT MOUSE REGIONAL RESOURCE CENTER: UC DAVIS
 2795 2nd Street, Suite 400, Davis, CA 95618
mmrrc@ucdavis.edu
 530-754-MMRRC

NAME OF PCR: B6;129-Kif3c^{tm2Gsn}/Mmcd MMRRC # 000215-UCD

Protocol:

Reagent/ Constituent	Volume (μL)
Water	15.4
10x Buffer (contains / without 15mM MgCl ₂)	2.5
dNTPs (stock concentration is 10mM)	0.5
Primer 1 (stock concentration is 20μM) B51820	0.2
Primer 2 (stock concentration is 20μM) B5RVR	0.2
Taq Polymerase 5Units/μL	0.2
DNA extracted with ☒ NaOH ☐ Proteinase K ☐ Other:	1.0
TOTAL VOLUME OF REACTION:	20μL

Comments on protocol:

Strategy:

Steps	Temp (°C)	Time (m:ss)	# of Cycles
1. Initiation/Melting HOT START? ☐	94	5:00	1
2. Denaturation	94	0:45	} 35x
3. Annealing	60	0:45	
4. Elongation	72	1:30	
5. Amplification	72	10:00	1
6. Finish	4	∞	n/a


Primers:

Name	Nucleotide Sequence (5' - 3')
1: B51820	GGA GAA TTA CCT GCA GGA CC
2: B5RVR	TGT AAG TGC ACC TAG CTA CC

Electrophoresis Protocol:

Agarose: 2% V: 100
 Estimated Running Time: 60 min.

Expected Bands	Genotype
350 bp	WT +/+
350 / 450 bp	HET +/-
450 bp	KO -/-



Lanes:
 1 Kb+ ladder
 2. H2O
 3. B6
 4. Kif3c null type II control
 5+. Samples