GENOTYPING BY PCR PROTOCOL MUTANT MOUSE REGIONAL RESOURCE CENTER: UC DAVIS

2795 2nd Street, Suite 400, Davis, CA 95618

mmrrc@ucdavis.edu

530-754-MMRRC

NAME OF PCR: C57BL/6-Lyst^{bg-Btlr}/Mmcd, (souris) MMRRC # 010470-UCD

Protocol:

Reagent/ Constituent	Volume (µL)
Water	12.675
10x Buffer (contains 15mM MgCl ₂)	2.5
Betaine (stock concentration is 5M) Optional	6.5
dNTPs (stock concentration is 25mM)	0.5
DMSO Optional	0.325
Primer 1 (stock concentration is 20µM) Souris PCR F	0.5
Primer 2 (stock concentration is 20µM) Souris PCR R	0.5
Taq Polymerase	0.5
DNA sample extracted with ☐ NaOH ☐ Proteinase K ☐ Other: Any	1.0
TOTAL VOLUME OF REACTION:	25µL

Comments on protocol:

- PCR products are verified to contain the correct amplicon size by running ~10µl of the reaction on a gel and the remaining 15µl purified via column based PCR purification method for sequencing.
- Use Touch-Down cycling protocol-first 10 cycles anneal at 65° C decreasing in temperature by 1.0° C; next 30 cycles anneal at 55° C.
- Betaine and DMSO have been standardized due to high GC content. Protocol may be tested without. Also, may adjust MgCl₂ to increase reaction or decrease non specific amplifications.

Strategy:

Steps	Temp (°C)	Time (m:ss)	# of Cycles
1. Initiation/Melting HOT START?	94	5:00	1
2. Denaturation	94	0:15	
3. Annealing steps 2-3-4 will cycle in sequence	65 to 55 (\1°C/cycle)	0:30	1 0x
4. Elongation	72	0:40	J
5. Denaturation	94	0:15	
6. Annealing steps 5-6-7 will cycle in sequence	55	0:30	30x
7. Elongation	72	0:40	J
8. Amplification	72	5:00	1
9. Finish	15	∞	n/a

Primers:

Name	Nucleotide Sequence (5' - 3')
1: Souris PCR F	TCA CCC TTC ATG TAG ACC AGG AAC C
2: Souris PCR R	TGC AGG GCC AAA GCC ATA TCT AAA C
3: Souris _seq F	CAG AAG TCA ATG TGC CTT ACT TGC
4: Souris _seq R	GAT GAC CTT GAA ATT CTG ACC ATC C

Electrophoresis Protocol:

Agarose: 2% mV: 80 Estimated Running Time: 90 min

Primer Combination	Band	Genotype
1 and 2	bp	souris

Mutation site (red) and flanking sequence:

WT aaagTaagaa souris aaagAaagaa