

GENOTYPING BY PCR PROTOCOL

MUTANT MOUSE REGIONAL RESOURCE CENTER: UC DAVIS

mmrrc@ucdavis.edu
530-754-MMRRC

NAME OF PCR: **B6;129P2-*Hvcn1*^{Gt(RRN293)Byg}/Mmucd** MMRRC: **015990-UCD**

Protocol:

Reagent/Constituent	Volume (μL)
Water	10.275
10x Buffer	2.5
MgCl ₂ (stock concentration is 25mM)	1.7
Betaine (stock concentration is 5M) <i>Optional</i>	6.5
dNTPs (stock concentration is 10mM)	0.5
DMSO <i>Optional</i>	0.325
Primer 1. (stock concentration is 20μM) β-Geo F	0.5
Primer 2. (stock concentration is 20μM) β-Geo R	0.5
Primer 3. (stock concentration is 20μM) Tcrd F	0.5
Primer 4. (stock concentration is 20μM) Tcrd R	0.5
Taq Polymerase 5Units/μL	0.2
DNA (example) extracted w/ Proteinase K	1.0
TOTAL VOLUME OF REACTION:	25.000 μL

Comments on protocol:

- Protocol may work with other DNA extraction methods.
- Use Touch-Down cycling protocol-first 10 cycles anneal at 65°C decreasing in temperature by 1.0°C; next 30 cycles anneal at 55°C.
- Betaine and DMSO have been standardized due to high GC content. Protocol may be tested without. Also, may adjust MgCl₂ to increase reaction or decrease non-specific amplifications.

Strategy:

Steps	Temp (°C)	Time (m:ss)	# of Cycles
1. Initiation/Melting HOT START? ☐	94	5:00	1
2. Denaturation	94	0:15	
3. Annealing steps 2-3-4 cycle in sequence	65 to 55 (↓1°C/cycle)	0:30	40x
4. Elongation	72	0:40	
5. Amplification	72	5:00	1
6. Finish	4	∞	n/a

Primers:

Electrophoresis Protocol:

Name	Nucleotide Sequence (5' - 3')	Agarose: 1.5% V: 90		
1. β-Geo F	CAAATGGCGATTACCGTTGA	Estimated Running Time: 90 min.		
2. β-Geo R	TGCCAGTCATAGCCGAATA	Primer Combination	Band (bp)	Genotype
3. Tcrd F	CAAATGTTGCTTGCTGCTGGTG	1 and 2	581	β-Geo +
4. Tcrd R	GTCAGTCGAGTGCACAGTTT	3 and 4	200	DNA ref rxn

