

GENOTYPING BY PCR PROTOCOL FORM

MUTANT MOUSE REGIONAL RESOURCE CENTER: UC DAVIS

Investigator/PI: Bruce Beutler, M.D. Address: N. Torrey Pines Rd., SP293 Contact: Xin Du

Telephone: 858-784-8610 FAX: 858-784-8444 email: bruce@scripps.edu

DNA Extraction Method: NaOH _____ Proteinase K X Other _____

Protocol: **NAME OF PCR:** MMRRC Strain #30754-UCD, C57BL/6J-Tmprss6^{m2Btlr} (Zorro)

Reagent/ Constituent	Volume (uL)
DNA Sample	1
10x Buffer (contains 15mM MgCl ₂)	5
dNTPs (stock concentration is 25mM)	0.4
Primer 1 (stock concentration is 20 uM)	1
Primer 2 (stock concentration is 20 uM)	1
Primer 3 (stock concentration is 20 uM)	
Primer 4 (stock concentration is 20 uM)	
Taq Polymerase	2.5
water	39.1
TOTAL VOLUME OF REACTION:	
	50 uL

Comments on protocol (e.g., different concentration of MgCl₂, etc): using primers 1 & 2 for amplification and 3 & 4 for sequencing.

Strategy:

Steps	Temp (°C)	Time (min)	# of Cycles
1. Initiation/Melting HOT START?..CHECK HERE []	94	2	1
2. Denaturation	94	30 seconds	30
3. Annealing } steps 2-3-4 will cycle in sequence	55	30 seconds	
4. Elongation	72	1 min	
5. Amplification (i.e., 72°C, 10 min)	72	7	1
6. Finish (i.e., 4°C, indefinite)	4	n/a	n/a

Primers:

Primer Name	Nucleotide Sequence (5' - 3')
1:TmpRSS6_zorroF	CAGTACAGCCATTGGCATGTAGGAG
2:TmpRSS6_zorroR	TGTGTGAAGAAGCCCAACCCAGAG
3: zorro_seqF	CTAGCAACACACTGCATGTTTG
4: zorro_seqR	TCAGATTGCAGAGACGGCTC

Electrophoresis Protocol:

% Agarose: _____ V : _____

Estimated Running Time (min): _____

Primer combination	Band (bp)	genotype
(i.e. 1&2)	769	
(i.e. 3&4)		
(i.e. 1&2&3)		

The flanking sequences of mutation sites:

Wt CGGGGT**C**GACACA
 zorro CGGGGT**T**GACACA