

# GENOTYPING BY PCR PROTOCOL

## MUTANT MOUSE REGIONAL RESOURCE CENTER: UC DAVIS

[mmrrc@ucdavis.edu](mailto:mmrrc@ucdavis.edu)  
530-754-MMRRC

NAME OF PCR: FVB-Tg(TSPY1-Figla)1Dean/Mmcd MMRRC: 032870-UCD

### Protocol:

| Reagent/Constituent                                                                   | Volume (μL)      |
|---------------------------------------------------------------------------------------|------------------|
| Water                                                                                 | 11.275           |
| 10x Buffer                                                                            | 2.5              |
| MgCl <sub>2</sub> (stock concentration is 25mM)                                       | 1.7              |
| Betaine (stock concentration is 5M) <i>Optional</i>                                   | 6.5              |
| dNTPs (stock concentration is 10mM)                                                   | 0.5              |
| DMSO <i>Optional</i>                                                                  | 0.325            |
| Primer 1. (stock concentration is 20μM)                                               | 0.5              |
| Primer 2. (stock concentration is 20μM)                                               | 0.5              |
| Taq Polymerase 5Units/μL                                                              | 0.2              |
| DNA (example) extracted w/ "Qiagen DNeasy columns or other similar silica based kits" | 1.0              |
| <b>TOTAL VOLUME OF REACTION:</b>                                                      | <b>25.000 μL</b> |

### Comments on protocol:

- Protocol may work with other DNA extraction methods.
- Use Touch-Down cycling protocol-first 10 cycles anneal at 65°C decreasing in temperature by 1.0°C; next 30 cycles anneal at 55°C.
- Betaine and DMSO have been standardized due to high GC content. Protocol may be tested without. Also, may adjust

### Strategy:

| Steps                                                                                        | Temp (°C)             | Time (m:ss) | # of Cycles |
|----------------------------------------------------------------------------------------------|-----------------------|-------------|-------------|
| 1. Initiation/Melting <span style="float: right;">HOT START? <input type="checkbox"/></span> | 94                    | 5:00        | <b>1</b>    |
| 2. Denaturation                                                                              | 94                    | 0:15        |             |
| 3. Annealing <span style="float: right;">steps 2-3-4 cycle in sequence</span>                | 65 to 55 (↓1°C/cycle) | 0:30        | <b>40x</b>  |
| 4. Elongation                                                                                | 72                    | 0:40        |             |
| 5. Amplification                                                                             | 72                    | 5:00        | <b>1</b>    |
| 6. Finish                                                                                    | 15                    | ∞           | n/a         |

### Primers:

| Name            | Nucleotide Sequence (5' - 3') | Electrophoresis Protocol:              |                  |                 |
|-----------------|-------------------------------|----------------------------------------|------------------|-----------------|
| 1. FWD TSPY2314 | CACCGCCCTCCCATTGACTGGCCACGAC  | Argarose: <b>1.5%</b>                  | V: <b>90</b>     |                 |
| 2. Rev FIG180R  | GCGCTTGAGCCTACAGATGGCGGCGAGCC | Estimated Running Time: <b>90</b> min. |                  |                 |
|                 |                               | <b>Primer Combination</b>              | <b>Band (bp)</b> | <b>Genotype</b> |
|                 |                               | 1 and 2                                | 252 bp           | +               |

