

GENOTYPING BY PCR PROTOCOL

MUTANT MOUSE RESOURCE & RESEARCH CENTER: UC DAVIS

mmrrc@ucdavis.edu

530-754-MMRRC

Protocol Name: C57BL/6N-Rhot1^{tm1a(EUCOMM)Wtsi}/Mmucd MMRRC: 037029-UCD

Protocol:

Reagent/Constituent	Volume (μL)
Water	11.275
10x Buffer	2.5
MgCl ₂ (stock concentration is 25mM)	1.7
Betaine (stock concentration is 5M) <i>Optional</i>	6.5
dNTPs (stock concentration is 10mM)	0.5
DMSO <i>Optional</i>	0.325
Primer 1. (stock concentration is 20μM)	0.5
Primer 2. (stock concentration is 20μM)	0.5
Taq Polymerase 5Units/μL	0.2
DNA (example) extracted w/ "Qiagen DNeasy columns or other similar silica based kits"	1.0
TOTAL VOLUME OF REACTION:	25.000 μL

Comments on protocol:

- Protocol may work with other DNA extraction methods.
- Use Touch-Down cycling protocol-first 10 cycles anneal at 65°C decreasing in temperature by 1.0°C; next 30 cycles anneal at 55°C.
- Betaine and DMSO have been standardized due to high GC content. Protocol may be tested without. Also, may adjust MgCl₂ to increase reaction or decrease non-specific amplifications.

Strategy:

Steps	Temp (°C)	Time (m:ss)	# of Cycles
1. Initiation/Melting HOT START? ☐	94	5:00	1
2. Denaturation	94	0:15	
3. Annealing steps 2-3-4 cycle in sequence	65 to 55 (↓1°C/cycle)	0:30	40x
4. Elongation	72	0:40	
5. Amplification	72	5:00	1
6. Finish	15	∞	n/a

Primers:

Name	Nucleotide Sequence (5' - 3')
1. 37029-lacF	GCTACCATTACCAGTTGGTCTGGTG
2. 37029-neoF	GGGATCTCATGCTGGAGTTCTTCG
3. 37029-loxF	GAGATGGCGCAACGCAATTAATG
4. 37029-TTR	TGGTAAGCCAGTATTCCCAGCTGTAG
5. 37029-R	AGCACATCACCACTGCCTTCC
6. 37029-F	TGAGCTCTGCTCAGTGCAGTGG

Electrophoresis Protocol:

Argarose:	1.5%	V:	90
Estimated Running Time:	90 min.		
Primer Combination	Band (bp)	Genotype	
3 & 5	202	Floxed	
2 & 4	593	Pre-Cre	
1 & 5	549	Post-Cre	
4 & 6	497	Wild-type	
4 & 6	543	Post-Flp	
5 & 6	499	Post-Flp & Post-Cre	

