

KOMP PCR Design

Mouse PCR Protocol (version v2JE)

Design ID: 43838

Project ID: CSD34319

Selection Cassette: L1L2_Bact_P



MMRRC Stock #: 046367-UCD

C57BL/6N-4921536K21Riktm1a(KOMP)Wtsi/
Mmucd


Suggested DNA Prep: DNeasy®Tissue Kit

Reagent	1X (μL)
water (biological grade)	10.725
betain 5M (Sigma)	6.5
DMSO (Sigma)	0.325
10X buffer w/o MgCl ₂ (AB)	2.5
25 mM MgCl ₂ (AB)	1.75
10 mM dNTPs (Invitrogen)	0.5
primers (20 μM each)	0.5
Taq 5U/μL (AmpliTaq, AB)	0.2
total cocktail	23
template	2
reaction volume	25

Cycling Parameters

Temperature °C	Time	
94	5 min	
94	15 sec	10X (decrease 1°C/cycle)
65	30 sec	
72	40 sec	
94	15 sec	
55	30 sec	30X
72	40 sec	
72	5 min	
4	finished	

Primer Strategy



Cassette Primers

CSD-lacF: GCTACCATTACCAGTTGGTCTGGTGTC
CSD-neoF: GGGATCTCATGCTGGAGTTCTTCG
CSD-loxF: GAGATGGCGCAACGCAATTAATG

Gene Specific Primers

CSD-4921536K21Rik-R: CTGGTCTATCTCTGAGTCCCAACCC
CSD-4921536K21Rik-ttR: GCAAGCCAGTAAAGAACATCCCTCC
CSD-4921536K21Rik-F: CTGAAAATGAGAAAACCACTTCTCTGG

Genotype	Forward Primer	Reverse Primer	Amplicon size (bp)
Floxed	CSD-loxF	CSD-4921536K21Rik-R	288
PreCre	CSD-neoF	CSD-4921536K21Rik-ttR	567
PostCre	CSD-lacF	CSD-4921536K21Rik-R	586
Wildtype	CSD-4921536K21Rik-F	CSD-4921536K21Rik-ttR	304
PostFlp	CSD-4921536K21Rik-F	CSD-4921536K21Rik-ttR	449
PostFlp & Cre	CSD-4921536K21Rik-F	CSD-4921536K21Rik-R	517

Please note, these primers are auto-designed and may not have been verified by the repository, and as such may require optimization or redesign by your facility.

We recommend running primers singleplex. For screening of pups prior to any Cre recombination, the PreCre primers may be used to identify the mutant mice. The PostCre primers should be used if mutant mice were crossed with a Cre recombinase line. The wildtype primers should be used for zygosity testing of all mutant mice (pre or post recombination).