

# KOMP PCR Design

Mouse PCR Protocol (version 1)

Design ID: 42399

Project ID: CSD37393

Selection Cassette: L1L2\_gt2

MMRRC Stock #: 047139-UCD

C57BL/6N-Cluap1tm1a(KOMP)Wtsi/Mmucd


## Suggested DNA Prep: DNeasy® Tissue Kit

Reagent	1X ( μL)
water (biological grade)	10.725
betain 5M (Sigma)	6.5
DMSO (Sigma)	0.325
10X buffer w/o MgCl <sub>2</sub> (AB)	2.5
25 mM MgCl <sub>2</sub> (AB)	1.75
10 mM dNTPs (Invitrogen)	0.5
primers (20 μM each)	0.5
Taq 5U/ μL (AmpliTaq, AB)	0.2
total cocktail	23
template	2
reaction volume	25

### Cycling Parameters

Temperature °C	Time	
94	5 min	
94	15 sec	<b>10X (decrease 1°C/cycle)</b>
65	30 sec	
72	40 sec	
94	15 sec	<b>30X</b>
55	30 sec	
72	40 sec	
72	5 min	
4	finished	

## Primer Strategy



### Cassette Primers

CSD-neoF: GGGATCTCATGCTGGAGTTCTTCG  
CSD-loxF: GAGATGGCGCAACGCAATTAATG

### Gene Specific Primers

CSD-Cluap1-R: CTGGCTTCTAAGAATAAGCTGGCCG  
CSD-Cluap1-ttR: ATTCCAACCCACACCTGAGATTCCG  
CSD-Cluap1-F: GTAATTTCCACAGCTCTGCCACTCC

Genotype	Forward Primer	Reverse Primer	Amplicon size (bp)
Floxed	CSD-loxF	CSD-Cluap1-R	338
PreCre	CSD-neoF	CSD-Cluap1-ttR	517
PostCre	CSD-neoF	CSD-Cluap1-R	685
Wildtype	CSD-Cluap1-F	CSD-Cluap1-ttR	506
PostFlp	CSD-Cluap1-F	CSD-Cluap1-ttR	659
PostFlp & Cre	CSD-Cluap1-F	CSD-Cluap1-R	827

Please note, these primers are auto-designed and may not have been verified by the repository, and as such may require optimization or redesign by your facility.

We recommend running primers singleplex. For screening of pups prior to any Flp or Cre recombination, the Floxed or PreCre primers may be used to identify the mutant mice. The Floxed primers test for the distal LoxP site. The PostCre primers should be used if mutant mice were crossed with a Cre recombinase line (without any FLP recombination). The PostFlp primers should be used if mutant mice were crossed with a Flp recombinase line. The PostFlp & Cre primers should be used if mutant mice were crossed with a Flp recombinase line and then a Cre recombinase line. The wildtype primers should be used for zygosity testing of all mutant mice (pre or post recombination).