

KOMP PCR Design

Mouse PCR Protocol (version 2BW)

Design ID: 49236

Project ID: CSD40214

Selection Cassette: L1L2_Bact_P




MMRRC Stock #: 047310-UCD

C57BL/6N-Dbn1tm1b(KOMP)Wtsi/JMmucd

Suggested DNA Prep: DNeasy®Tissue Kit

Reagent	1X (μL)	Cycling Parameters		
water (biological grade)	10.725	Temperature °C	Time	
betain 5M (Sigma)	6.5	94	5 min	
DMSO (Sigma)	0.325	94	15 sec	10X (decrease 1°C/cycle)
10X buffer w/o MgCl ₂ (AB)	2.5	65	30 sec	
25 mM MgCl ₂ (AB)	1.75	72	40 sec	
10 mM dNTPs (Invitrogen)	0.5	94	15 sec	30X
primers (20 μM each)	0.5	55	30 sec	
Taq 5U/μL (AmpliTaQ, AB)	0.2	72	40 sec	
total cocktail	23	72	5 min	
template	2	4	finished	
reaction volume	25			

Primer Strategy



Cassette Primers

CSD-lacF: GCTACCATTACCAGTTGGTCTGGTGTC
 CSD-neoF: GGGATCTCATGCTGGAGTTCTTCG
 CSD-loxF: GAGATGGCGCAACGCAATTAATG

Gene Specific Primers

CSD-Dbn1-R: AGAGCCTGTCTATGTGTTTCAGTCCC
 CSD-Dbn1-ttR: ATCTCCATTTCTCGGGGATCACACC
 CSD-Dbn1-F2: GTTTTGTCCCTCTCGCTTTCC

Genotype	Forward Primer	Reverse Primer	Amplicon size (bp)
Floxed	CSD-loxF	CSD-Dbn1-R	236
PreCre	CSD-neoF	CSD-Dbn1-ttR	668
PostCre	CSD-lacF	CSD-Dbn1-R	534
Wildtype	CSD-Dbn1-F	CSD-Dbn1-ttR	322
PostFlp	CSD-Dbn1-F	CSD-Dbn1-ttR	509
PostFlp & Cre	CSD-Dbn1-F	CSD-Dbn1-R	424

Please note, these primers are auto-designed and may not have been verified by the repository, and as such may require optimization or redesign by your facility.

We recommend running primers singleplex. For screening of pups prior to any Flp or Cre recombination, the Floxed or PreCre primers may be used to identify the mutant mice. The Floxed primers test for the distal LoxP site. The PostCre primers should be used if mutant mice were crossed with a Cre recombinase line (without any FLP recombination). The PostFlp primers should be used if mutant mice were crossed with a Flp recombinase line. The PostFlp & Cre primers should be used if mutant mice were crossed with a Flp recombinase line and then a Cre recombinase line. The wildtype primers should be used for zygosity testing of all mutant mice (pre or post recombination).