

# KOMP PCR Design

Mouse PCR Protocol (version 1JE)

Design ID: 45655

Project ID: CSD79901

Selection Cassette: L1L2\_Bact\_P

**MMRRC @ UC Davis**  
Mutant Mouse Resource & Research Centers supported by NIH

MMRRC Stock #: 049511-UCD

C57BL/6N-Atm1Brd Rrm2tm1a(KOMP)Wtsi/  
Mmucd


## Suggested DNA Prep: DNeasy®Tissue Kit

Reagent	1X (µL)
water (biological grade)	10.725
betain 5M (Sigma)	6.5
DMSO (Sigma)	0.325
10X buffer w/o MgCl <sub>2</sub> (AB)	2.5
25 mM MgCl <sub>2</sub> (AB)	1.75
10 mM dNTPs (Invitrogen)	0.5
primers (20 µM each)	0.5
Taq 5U/µL (AmpliAq, AB)	0.2
total cocktail	23
template	2
reaction volume	25

### Cycling Parameters

Temperature °C	Time	
94	5 min	
94	15 sec	<b>10X (decrease 1°C/cycle)</b>
65	30 sec	
72	40 sec	
94	15 sec	<b>30X</b>
55	30 sec	
72	40 sec	
72	5 min	
4	finished	

## Primer Strategy



### Cassette Primers

CSD-lacF: GCTACCATTACCAGTTGGTCTGGTGTC  
 CSD-neoF: GGGATCTCATGCTGGAGTTCTTCG  
 CSD-loxF: GAGATGGCGCAACGCAATTAATG

### Gene Specific Primers

CSD-R: CGTGTCAGACACCTTATGTGTGATGC  
 CSD-ttR: GTTCTTCATGATTTGCAAGCAAGC  
 CSD-F: ATACATGCACGGACCCATTCCC

### Genotype Forward Primer Reverse Primer Amplicon size (bp)

Genotype	Forward Primer	Reverse Primer	Amplicon size (bp)
Floxed	CSD-loxF	CSD-R	271
PreCre	CSD-neoF	CSD-ttR	559
PostCre	CSD-lacF	CSD-R	569
Wildtype	CSD-F	CSD-ttR	397
PostFlp	CSD-F	CSD-ttR	492
PostFlp & Cre	CSD-F	CSD-R	551

Please note, these primers are auto-designed and may not have been verified by the repository, and as such may require optimization or redesign by your facility.

We recommend running primers singleplex. For screening of pups prior to any Flp or Cre recombination, the Floxed or PreCre primers may be used to identify the mutant mice. The Floxed primers test for the distal LoxP site. The PostCre primers should be used if mutant mice were crossed with a Cre recombinase line (without any FLP recombination). The PostFlp primers should be used if mutant mice were crossed with a Flp recombinase line. The PostFlp & Cre primers should be used if mutant mice were crossed with a Flp recombinase line and then a Cre recombinase line. The wildtype primers should be used for zygosity testing of all mutant mice (pre or post recombination).