

KOMP PCR Design

Mouse PCR Protocol (version 2BW)

Design ID: 220971

Project ID: CSD26838

Selection Cassette: L1L2_Bact_P



MMRRC Stock #: 049744-UCD

C57BL/6N-Slc52a2tm1.1(KOMP)Vlclg/MbpMmucd


Suggested DNA Prep: DNeasy®Tissue Kit

Reagent	1X (μL)
water (biological grade)	10.725
betain 5M (Sigma)	6.5
DMSO (Sigma)	0.325
10X buffer w/o MgCl ₂ (AB)	2.5
25 mM MgCl ₂ (AB)	1.75
10 mM dNTPs (Invitrogen)	0.5
primers (20 μM each)	0.5
Taq 5U/μL (AmpliTaq, AB)	0.2
total cocktail	23
template	2
reaction volume	25

Cycling Parameters

Temperature °C	Time	
94	5 min	
94	15 sec	10X (decrease 1°C/cycle)
65	30 sec	
72	40 sec	
94	15 sec	
55	30 sec	30X
72	40 sec	
72	5 min	
4	finished	

Primer Strategy



Cassette Primers

CSD-lacF: GCTACCATTACCAAGTTGGTCTGGTGTGTC
CSD-neoF: GGGATCTCATGCTGGAGTTCTTCG

Gene Specific Primers

CSD-Slc52a2-R: TGGTCCTCACACTGGTCTTTTCTCC
CSD-Slc52a2-F: CCCGCTTTGACCTTTGACCTAATGG
CSD-Slc52-wtR: CCAGCAGCAGACCCAGGTTTCC

Genotype Forward Primer Reverse Primer Amplicon size (bp)

PreCre	CSD-neoF	CSD-Slc52a2-R	629
PostCre	CSD-lacF	CSD-Slc52a2-R	580
Wildtype	CSD-Slc52a2-F	CSD-Slc52a2-wtR	305
PostFlp	CSD-Slc52a2-F	CSD-Slc52a2-R	611

Please note, these primers are auto-designed and may not have been verified by the repository, and as such may require optimization or redesign by your facility.

We recommend running primers singleplex. For screening of pups prior to any Cre recombination, the PreCre primers may be used to identify the mutant mice. The PostCre primers should be used if mutant mice were crossed with a Cre recombinase line. The wildtype primers should be used for zygosity testing of all mutant mice (pre or post recombination).