

GENOTYPING BY PCR PROTOCOL

MUTANT MOUSE RESOURCE & RESEARCH CENTER: UC DAVIS

mmrrc@ucdavis.edu

530-754-MMRRC

Protocol Name: C57BL/6N-Abatem1(IMPC)Bay/Mmucd **MMRRC: 050975-UCD**

Protocol:

Reagent/Constituent	Volume (μL)
Water	5.6
GoTaq® G2 Colorless Master Mix, 2X	7.5
Primer 1. (stock concentration is 20μM)	0.45
Primer 2. (stock concentration is 20μM)	0.45
DNA (example) extracted w/ "Qiagen DNeasy columns or other similar silica based kits"	1.0
TOTAL VOLUME	
15	

Comments on protocol:

- Protocol may work with other DNA extraction methods.
- Use Touch-Down cycling protocol-first 10 cycles anneal at 65°C decreasing in temperature by 1.0°C; next 30 cycles anneal at 55°C.

Strategy:

Steps	Temp (°C)	Time (m:ss)	# of Cycles
1. Initiation/Melting HOT START? ☐	94	2:00	1x
2. Denaturation	94	0:10	
3. Annealing steps 2-3-4 cycle in sequence	65 (↓1°C/cycle)	0:30	10x
4. Elongation	68	2:00	
5. Denaturation	94	0:15	
6. Annealing steps 5-6-7 cycle in sequence	55	0:30	25x
7. Elongation	68	2:00 (↑20sec/cycle)	

Primers:

Electrophoresis Protocol:

Name	Nucleotide Sequence (5' - 3')	Argarose: 1.5% V: 90		
1. 50975-5LoxPF	CTGGAAGAGCAGCTTTTGGC	Estimated Running Time: 90 min.		
2. 50975-5LoxPR	AGCTCTCCTCCCAGTCCAAA	Primer Combination	Band (bp)	Genotype
3. 50975-3LoxPF	AACAAAGACCCAGGAGGCAG	1 & 2	219, 179	floxed, wildtype
4. 50975-3LoxPR	CACAATCTGGCTGCCACATG	3 & 4	221, 181	floxed, wildtype
5. 50975-KOF	GCCAAAGTTGAGAGACTCCT	5 & 6	407, 1114	KO, wildtype
6. 50975-KOR	CACCGACCTAACTGCTGGTT	5 & 7	552	wildtype
7. 50975-EndoR	CTCCTGTCCTAAGGCAGCAG			

GENOTYPING BY PCR PROTOCOL
MUTANT MOUSE RESOURCE & RESEARCH CENTER: UC DAVIS

mmrrc@ucdavis.edu
530-754-MMRRC

