

# GENOTYPING PROTOCOL

## MUTANT MOUSE RESOURCE & RESEARCH CENTER: UC DAVIS

[mmrrc@ucdavis.edu](mailto:mmrrc@ucdavis.edu)

530-754-MMRRC

Protocol Name: CR11422 AI593442 EXDEL

Protocol: GoTaq® G2 Colorless Master Mix(Promega)

| Reagent/Constituent                                                                   | Volume (μL)     |
|---------------------------------------------------------------------------------------|-----------------|
| Water                                                                                 | 4.5             |
| GoTaq® G2 Colorless Master Mix,2X                                                     | 7.5             |
| Primer 1. (stock concentration is 20μM) comF                                          | 0.5             |
| Primer 2. (stock concentration is 20μM) wtR                                           | 0.5             |
| Primer 3. (stock concentration is 20μM) mutR                                          | 0.5             |
| DNA (example) extracted w/ "Qiagen DNeasy columns or other similar silica based kits" | 1.5             |
| <b>TOTAL VOLUME OF REACTION:</b>                                                      | <b>15.00 μL</b> |

### Comments on protocol:

- Protocol may work with other DNA extraction methods.

### Strategy:

| Steps                                                     | Temp (°C )      | Time (m:ss)         | # of Cycles |
|-----------------------------------------------------------|-----------------|---------------------|-------------|
| 1. Initiation/Melting HOT START? <input type="checkbox"/> | 94              | 2:00                | 1x          |
| 2. Denaturation                                           | 94              | 0:10                |             |
| 3. Annealing steps 2-3-4 cycle in sequence                | 65 (↓1°C/cycle) | 0:30                | 10x         |
| 4. Elongation                                             | 68              | 2:00                |             |
| 5. Denaturation                                           | 94              | 0:15                |             |
| 6. Annealing steps 5-6-7 cycle in sequence                | 55              | 0:30                | 25x         |
| 7. Elongation                                             | 68              | 2:00 (↑20sec/cycle) |             |
| 8. Finish                                                 | 4               | ∞                   | n/a         |

### Primers:

| Name                | Nucleotide Sequence (5' - 3') | Agarose: 1.5% V: 90                                        |
|---------------------|-------------------------------|------------------------------------------------------------|
| 1. CR_AI593442_comF | TTCTTTGCAGCACTGTGCCT          | Estimated Running Time: 90 min.                            |
| 2. CR_AI593442_wtR* | GTTGTGGGAAGCAAAGGTCCG         | <b>Primer Combination</b> <b>Band (bp)</b> <b>Genotype</b> |
| 3. CR_AI593442_mutR | TGCAGCCTTCATTAACAATGAGA       | 1 & 2, 1 & 3 330, 1169 wildtype                            |
|                     |                               | 1 & 3 305 mutant                                           |

### Electrophoresis Protocol:

**Allele Description:** Exon 2 ENSMUSE00001397118 was constitutively deleted from the 5'UTR through the 3' UTR from the AI593442 Gene ENSMUSG00000078307 using CRISPR Cas9 gene editing technology in mouse zygotes. The subsequent 864bp deletion from Chr 9: 52,588,824-52,589,687 GRCm39 was screened by PCR analysis. The selected founder animal was backcrossed to C57BL6/N to produce sequence confirmed heterozygous animals for establishing and archiving the line.

\*wtR primer untested (ePCR verified)

