# GENOTYPING PROTOCOL MUTANT MOUSE RESOURCE & RESEARCH CENTER: UC DAVIS

mmrrc@ucdavis.edu 530-754-MMRRC

Protocol Name: CR11429 Bri3bp EXDEL

Protocol: GoTaq® G2 Colorless Master Mix(Promega)

Reagent/Constituent	Volume (µL)
Water	4.5
GoTaq® G2 Colorless Master Mix,2X	7.5
Primer 1. (stock concentration is 20µM) comF	0.5
Primer 2. (stock concentration is 20µM) wtR	0.5
Primer 3. (stock concentration is 20µM) mutR	0.5
DNA (example) extracted w/ "Qiagen DNeasy columns or other similar silica based kits"	1.5
TOTAL VOLUME OF REACTION:	15.00 µL

#### Comments on protocol:

Protocol may work with other DNA extraction methods.

### Strategy:

Steps		Temp (°C)	Time (m:ss)	# of Cycles	
1. Initiation/Melting	HOT START? ☐	94	2:00	1x	
2. Denaturation		94	0:10		
3. Annealing	steps 2-3-4 cycle in sequence	65 (↓1°C/cycle)	0:30	10x	
4. Elongation		68	2:00		
5. Denaturation		94	0:15		
6. Annealing	steps 5-6-7 cycle in sequence	55	0:30	25x	
7. Elongation		68	2:00 (†20sec/cycle)		
8. Finish		4	∞	n/a	

#### Primers:

## **Electrophoresis Protocol:**

Name	Nucleotide Sequence (5' - 3')	Agarose: 1.5% V:	90	
1. CR_Bri3bp_comF	CCAGATACCCTGACCAAGAGC	Estimated Running Time:	<b>90</b> min.	
2. CR_Bri3bp_wtR*	CGAGAAGACAGTAAGGAATCAAC	Primer Combination	Band (bp)	Genotype
3. CR_Bri3bp_mutR	AGGTGACACGACCTGATGTGA	1 & 2,1 & 3	377, 1364	wildtype
		1 & 3	347	mutant

Allele Description: Exon 2 ENSMUSE00000317891 and flanking splicing regions were constitutively deleted from the Bri3bp Gene ENSMUSG00000037905 using CRISPR Cas9 gene editing technology in mouse zygotes. The subsequent 1017bp deletion from Chr 5: 125,528,565 – 125,529,581 GRCm39 was screened by PCR analysis. The selected founder animal was backcrossed to C57BL6/N to produce sequence confirmed heterozygous animals for establishing and archiving the line.

\*wtR primer untested (ePCR verified)

