

# GENOTYPING PROTOCOL

## MUTANT MOUSE RESOURCE & RESEARCH CENTER: UC DAVIS

[mmrrc@ucdavis.edu](mailto:mmrrc@ucdavis.edu)

530-754-MMRRC

**Protocol Name:** CR11449 Mphosph6 EXDEL

**Protocol:** GoTaq® G2 Colorless Master Mix(Promega)

Reagent/Constituent	Volume (μL)
Water	4.5
GoTaq® G2 Colorless Master Mix,2X	7.5
Primer 1. (stock concentration is 20μM) comF	0.5
Primer 2. (stock concentration is 20μM) wtR	0.5
Primer 3. (stock concentration is 20μM) mutR	0.5
DNA (example) extracted w/ "Qiagen DNeasy columns or other similar silica based kits"	1.5
<b>TOTAL VOLUME OF REACTION:</b>	<b>15.00 μL</b>

**Comments on protocol:**

- Protocol may work with other DNA extraction methods.

**Strategy:**

Steps	Temp (°C )	Time (m:ss)	# of Cycles
1. Initiation/Melting HOT START? <input type="checkbox"/>	94	2:00	<b>1x</b>
2. Denaturation	94	0:10	
3. Annealing steps 2-3-4 cycle in sequence	65 (↓1°C/cycle)	0:30	<b>10x</b>
4. Elongation	68	2:00	
5. Denaturation	94	0:15	
6. Annealing steps 5-6-7 cycle in sequence	55	0:30	<b>25x</b>
7. Elongation	68	2:00 (↑20sec/cycle)	
8. Finish	4	∞	n/a

**Primers:**

**Electrophoresis Protocol:**

Name	Nucleotide Sequence (5' - 3')	Agarose: 1.5% V: 90
1. CR_Mphosph6_comF	GTTACTGATGGAGCTGGAGAGCT	Estimated Running Time: 90 min.
2. CR_Mphosph6_wtR*	ACGACTCCCTGAGTACCAGT	<b>Primer Combination</b> <b>Band (bp)</b> <b>Genotype</b>
3. CR_Mphosph6_mutR	CCACAGGTAGCTAAGATTTTCACAAG	1 & 2, 1 & 3 329, 1168 wildtype
		1 & 3 693 mutant

**Allele Description:** Exon 2 ENSMUSE00000320725 and flanking splicing regions were constitutively deleted from the Mphosph6 Gene ENSMUSG00000031843 using CRISPR Cas9 gene editing technology in mouse zygotes. The subsequent 475bp deletion from Chr 8: 118,525,641 – 118,526,115 GRCm39 was screened by PCR analysis. The selected founder animal was backcrossed to C57BL6/N to produce sequence confirmed heterozygous animals for establishing and archiving the line.

\*wtR primer untested (ePCR verified)

