

GENOTYPING PROTOCOL

MUTANT MOUSE RESOURCE & RESEARCH CENTER: UC DAVIS

mmrrc@ucdavis.edu

530-754-MMRRC

Protocol Name: CR11475 Khl38 EXDEL

Protocol: GoTaq® G2 Colorless Master Mix(Promega)

Reagent/Constituent	Volume (μL)
Water	4.5
GoTaq® G2 Colorless Master Mix,2X	7.5
Primer 1. (stock concentration is 20μM) comF	0.5
Primer 2. (stock concentration is 20μM) wtR	0.5
Primer 3. (stock concentration is 20μM) mutR	0.5
DNA (example) extracted w/ "Qiagen DNeasy columns or other similar silica based kits"	1.5
TOTAL VOLUME OF REACTION:	15.00 μL

Comments on protocol:

- Protocol may work with other DNA extraction methods.

Strategy:

Steps	Temp (°C)	Time (m:ss)	# of Cycles
1. Initiation/Melting HOT START? <input type="checkbox"/>	94	2:00	1x
2. Denaturation	94	0:10	
3. Annealing steps 2-3-4 cycle in sequence	65 (↓1°C/cycle)	0:30	10x
4. Elongation	68	2:00	
5. Denaturation	94	0:15	
6. Annealing steps 5-6-7 cycle in sequence	55	0:30	25x
7. Elongation	68	2:00 (↑20sec/cycle)	
8. Finish	4	∞	n/a

Primers:

Electrophoresis Protocol:

Name	Nucleotide Sequence (5' - 3')	Agarose: 1.5% V: 90
1. CR_Khl38_comF	TGGCAAATCTGATGAGAATCTGGAC	Estimated Running Time: 90 min.
2. CR_Khl38_wtR*	CCATCTGGTAACTCCTCATCCAT	Primer Combination Band (bp) Genotype
3. CR_Khl38_mutR	GCTTGTTACACGACTAAAGACAC	1 & 2, 1 & 3 401,1888 wildtype
		1 & 3 366 mutant

Allele Description: Exon 2 ENSMUSE00000390153 and flanking splicing regions were constitutively deleted from the Khl38 Gene ENSMUSG00000022357 using CRISPR Cas9 gene editing technology in mouse zygotes. The subsequent 1522bp deletion from Chr 15: 58185371-58186892 GRCm39 was screened by PCR analysis. The selected founder animal was backcrossed to C57BL6/N to produce sequence confirmed heterozygous animals for establishing and archiving the line.

*wtR primer untested (ePCR verified)

