

GENOTYPING PROTOCOL

MUTANT MOUSE RESOURCE & RESEARCH CENTER: UC DAVIS

mmrrc@ucdavis.edu

530-754-MMRRC

Protocol Name: CR11467 Ccdc97 EXDEL

Protocol: GoTaq® G2 Colorless Master Mix(Promega)

Reagent/Constituent	Volume (µL)
Water	4.5
GoTaq® G2 Colorless Master Mix,2X	7.5
Primer 1. (stock concentration is 20µM) comF	0.5
Primer 2. (stock concentration is 20µM) wtR	0.5
Primer 3. (stock concentration is 20µM) mutR	0.5
DNA (example) extracted w/ "Qiagen DNeasy columns or other similar silica based kits"	1.5
TOTAL VOLUME OF REACTION:	15.00 µL

Comments on protocol:

- Protocol may work with other DNA extraction methods.

Strategy:

Steps	Temp (°C)	Time (m:ss)	# of Cycles
1. Initiation/Melting HOT START? ☐	94	2:00	1x
2. Denaturation	94	0:10	
3. Annealing steps 2-3-4 cycle in sequence	65 (↓1°C/cycle)	0:30	10x
4. Elongation	68	2:00	
5. Denaturation	94	0:15	
6. Annealing steps 5-6-7 cycle in sequence	55	0:30	25x
7. Elongation	68	2:00 (↑20sec/cycle)	
8. Finish	4	∞	n/a

Primers:

Electrophoresis Protocol:

Name	Nucleotide Sequence (5' - 3')	Agarose: 1.5% V: 90
1. CR_Ccdc97_comF	TGTCCGACTGTCTTTCTACCTAGC	Estimated Running Time: 90 min.
2. CR_Ccdc97_wtR*	CGTCTGAGAGTCAAGAGGAGTCT	Primer Combination Band (bp) Genotype
3. CR_Ccdc97_mutR	GGACACAGTTCCCATGTCATTG	1 & 2,1 & 3 430, 2732 wildtype
		1 & 3 657 mutant

Allele Description: Exon 2-4 (ENSMUSE00001306292, ENSMUSE00001249827, ENSMUSE00001240780) and flanking splicing regions were constitutively deleted from the Ccdc97 Gene ENSMUSG00000002608 using CRISPR Cas9 gene editing technology in mouse zygotes. The subsequent 2075bp deletion from Chr 7: 25413705 - 25415779 GRCm39 was screened by PCR analysis. The selected founder animal was backcrossed to C57BL6/N to produce sequence confirmed heterozygous animals for establishing and archiving the line.

*wtR primer untested (ePCR verified)

