

GENOTYPING PROTOCOL

MUTANT MOUSE RESOURCE & RESEARCH CENTER: UC DAVIS

mmrrc@ucdavis.edu

530-754-MMRRC

Protocol Name: CR11481 Mrps18a EXDEL

Protocol: GoTaq® G2 Colorless Master Mix(Promega)

Reagent/Constituent	Volume (μL)
Water	4.5
GoTaq® G2 Colorless Master Mix,2X	7.5
Primer 1. (stock concentration is 20μM) comF	0.5
Primer 2. (stock concentration is 20μM) wtR	0.5
Primer 3. (stock concentration is 20μM) mutR	0.5
DNA (example) extracted w/ "Qiagen DNeasy columns or other similar silica based kits"	1.5
TOTAL VOLUME OF REACTION:	15.00 μL

Comments on protocol:

- Protocol may work with other DNA extraction methods.

Strategy:

Steps	Temp (°C)	Time (m:ss)	# of Cycles
1. Initiation/Melting HOT START? <input type="checkbox"/>	94	2:00	1x
2. Denaturation	94	0:10	
3. Annealing steps 2-3-4 cycle in sequence	65 (↓1°C/cycle)	0:30	10x
4. Elongation	68	2:00	
5. Denaturation	94	0:15	
6. Annealing steps 5-6-7 cycle in sequence	55	0:30	25x
7. Elongation	68	2:00 (↑20sec/cycle)	
8. Finish	4	∞	n/a

Primers:

Electrophoresis Protocol:

Name	Nucleotide Sequence (5' - 3')	Agarose: 1.5%	V: 90
1. CR_Mrps18a_comF	TGGTCAAAATCAAGATGAGTCTCAGC	Estimated Running Time: 90 min.	
2. CR_Mrps18a_wtR*	TGCAGCGATACTTACTACAGTCGT	Primer Combination	Band (bp)
3. CR_Mrps18a_mutR	GCTGTCTTAGGGTAGACTATATTTGGC	1 & 2, 1 & 3	438, 1168
		1 & 3	544
			Genotype
			wildtype
			mutant

Allele Description: Exon 2 ENSMUSE00001288864 and flanking splicing regions were constitutively deleted from the Mrps18a Gene ENSMUSG00000023967 using CRISPR Cas9 gene editing technology in mouse zygotes. The subsequent 624bp deletion from Chr 17: 46,428,703 -46,429,326 GRCh39 was screened by PCR analysis. The selected founder animal was backcrossed to C57BL6/N to produce sequence confirmed heterozygous animals for establishing and archiving the line.

*wtR primer untested (ePCR verified)

