

GENOTYPING PROTOCOL

MUTANT MOUSE RESOURCE & RESEARCH CENTER: UC DAVIS

mmrrc@ucdavis.edu

530-754-MMRRC

Protocol Name: CR11483 Ndubf7 EXDEL

Protocol: GoTaq® G2 Colorless Master Mix(Promega)

Reagent/Constituent	Volume (μL)
Water	4.5
GoTaq® G2 Colorless Master Mix,2X	7.5
Primer 1. (stock concentration is 20μM) comF	0.5
Primer 2. (stock concentration is 20μM) wtR	0.5
Primer 3. (stock concentration is 20μM) mutR	0.5
DNA (example) extracted w/ "Qiagen DNeasy columns or other similar silica based kits"	1.5
TOTAL VOLUME OF REACTION:	15.00 μL

Comments on protocol:

- Protocol may work with other DNA extraction methods.

Strategy:

Steps	Temp (°C)	Time (m:ss)	# of Cycles
1. Initiation/Melting HOT START? ☐	94	2:00	1x
2. Denaturation	94	0:10	
3. Annealing steps 2-3-4 cycle in sequence	65 (↓1°C/cycle)	0:30	10x
4. Elongation	68	2:00	
5. Denaturation	94	0:15	
6. Annealing steps 5-6-7 cycle in sequence	55	0:30	25x
7. Elongation	68	2:00 (↑20sec/cycle)	
8. Finish	4	∞	n/a

Primers:

Name	Nucleotide Sequence (5' - 3')	Agarose: 1.5%	V: 90
1. CR_Ndubf7_comF	TTGCCAGTGATAGCTGATCTACC	Estimated Running Time: 90 min.	
2. CR_Ndubf7_wtR*	CTCTGATGTCCCTTCATCTAGAGAT	Primer Combination	Band (bp)
3. CR_Ndubf7_mutR	CATCCTTGAAGTAGCACAGATGAAG	1 & 2, 1 & 3	392,1393
		1 & 3	406
			mutant

Electrophoresis Protocol:

Allele Description: Exon 2 ENSMUSE00000278001 and flanking splicing regions were constitutively deleted from the Ndubf7 Gene ENSMUSG00000033938 using CRISPR Cas9 gene editing technology in mouse zygotes. The subsequent 987bp deletion from Chr 8: 84,296,957-84,297,943 GRCm39 was screened by PCR analysis. The selected founder animal was backcrossed to C57BL6/N to produce sequence confirmed heterozygous animals for establishing and archiving the line.

*wtR primer untested (ePCR verified)

