

# GENOTYPING PROTOCOL

## MUTANT MOUSE RESOURCE & RESEARCH CENTER: UC DAVIS

[mmrrc@ucdavis.edu](mailto:mmrrc@ucdavis.edu)

530-754-MMRRC

**Protocol Name:** CR11505 Sf3b5 EXDEL

**Protocol:** GoTaq® G2 Colorless Master Mix(Promega)

Reagent/Constituent	Volume (μL)
Water	4.5
GoTaq® G2 Colorless Master Mix,2X	7.5
Primer 1. (stock concentration is 20μM) comF	0.5
Primer 2. (stock concentration is 20μM) wtR	0.5
Primer 3. (stock concentration is 20μM) mutR	0.5
DNA (example) extracted w/ "Qiagen DNeasy columns or other similar silica based kits"	1.5
<b>TOTAL VOLUME OF REACTION:</b>	<b>15.00 μL</b>

### Comments on protocol:

- Protocol may work with other DNA extraction methods.

### Strategy:

Steps	Temp (°C )	Time (m:ss)	# of Cycles
1. Initiation/Melting HOT START? <input type="checkbox"/>	94	2:00	1x
2. Denaturation	94	0:10	
3. Annealing steps 2-3-4 cycle in sequence	65 (↓1°C/cycle)	0:30	10x
4. Elongation	68	2:00	
5. Denaturation	94	0:15	
6. Annealing steps 5-6-7 cycle in sequence	55	0:30	25x
7. Elongation	68	2:00 (↑20sec/cycle)	
8. Finish	4	∞	n/a

### Primers:

Name	Nucleotide Sequence (5' - 3')	Agarose: 1.5% V: 90
1. CR_Sf3b5_comF	TGGACTACTTGTTCAGAAATGCAAC	Estimated Running Time: 90 min.
2. CR_Sf3b5_wtR*	AGGTCAGGACACAGACGAGAAG	<b>Primer Combination</b> <b>Band (bp)</b> <b>Genotype</b>
3. CR_Sf3b5_mutR	CTCTCACAAAACCTTGATCCACAGA	1 & 2, 1 & 3 320, 980 wildtype
		1 & 3 428 mutant

### Electrophoresis Protocol:

**Allele Description:** Exon 1 ENSMUSE00000665291 was constitutively deleted from the 5'UTR through the 3' UTR from the Sf3b5 Gene ENSMUSG00000078348 using CRISPR Cas9 gene editing technology in mouse zygotes. The subsequent 552bp deletion from Chr 10: 12884162-12884713 GRCm39 was screened by PCR analysis. The selected founder animal was backcrossed to C57BL6/N to produce sequence confirmed heterozygous animals for establishing and archiving the line.

\*wtR primer untested (ePCR verified)

