

GENOTYPING PROTOCOL

MUTANT MOUSE RESOURCE & RESEARCH CENTER: UC DAVIS

mmrrc@ucdavis.edu

530-754-MMRRC

Protocol Name: CR11517 Srp14 EXDEL

Protocol: GoTaq® G2 Colorless Master Mix(Promega)

Reagent/Constituent	Volume (μL)
Water	4.5
GoTaq® G2 Colorless Master Mix,2X	7.5
Primer 1. (stock concentration is 20μM) comF	0.5
Primer 2. (stock concentration is 20μM) wtR	0.5
Primer 3. (stock concentration is 20μM) mutR	0.5
DNA (example) extracted w/ "Qiagen DNeasy columns or other similar silica based kits"	1.5
TOTAL VOLUME OF REACTION:	15.00 μL

Comments on protocol:

- Protocol may work with other DNA extraction methods.

Strategy:

Steps	Temp (°C)	Time (m:ss)	# of Cycles
1. Initiation/Melting HOT START? ☐	94	2:00	1x
2. Denaturation	94	0:10	
3. Annealing steps 2-3-4 cycle in sequence	65 (↓1°C/cycle)	0:30	10x
4. Elongation	68	2:00	
5. Denaturation	94	0:15	
6. Annealing steps 5-6-7 cycle in sequence	55	0:30	25x
7. Elongation	68	2:00 (↑20sec/cycle)	
8. Finish	4	∞	n/a

Primers:

Electrophoresis Protocol:

Name	Nucleotide Sequence (5' - 3')	Agarose: 1.5% V: 90
1. CR_Srp14_comF	GTGGACTCTTGGGTGACTTGC	Estimated Running Time: 90 min.
2. CR_Srp14_wtR*	GGTCAGCTCCGTCAGGAAGT	Primer Combination Band (bp) Genotype
3. CR_Srp14_mutR	GTCCCAGACACACATAGTACGAGAA	1 & 2,1 & 3 477,4201 wildtype
		1 & 3 632 mutant

Allele Description: Exon 1-5 (ENSMUSE00000706401, ENSMUSE00001249362, ENSMUSE00001214858, ENSMUSE00001297352, ENSMUSE00000731125) were constitutively deleted from 5'UTR through the 3' UTR from the Srp14 Gene ENSMUSG00000009549 using CRISPR Cas9 gene editing technology in mouse zygotes. The subsequent 3569bp deletion from Chr 2: 118306607- 118310175 GRCm39 was screened by PCR analysis. The selected founder animal was backcrossed to C57BL6/N to produce sequence confirmed heterozygous animals for establishing and archiving the line.

*wtR primer untested (ePCR verified)

