

GENOTYPING PROTOCOL

MUTANT MOUSE RESOURCE & RESEARCH CENTER: UC DAVIS

mmrrc@ucdavis.edu

530-754-MMRRC

Protocol Name: CR11541 Bnpl EXDEL

Protocol: GoTaq® G2 Colorless Master Mix(Promega)

Reagent/Constituent	Volume (μL)
Water	4.5
GoTaq® G2 Colorless Master Mix,2X	7.5
Primer 1. (stock concentration is 20μM) comF	0.5
Primer 2. (stock concentration is 20μM) wtR	0.5
Primer 3. (stock concentration is 20μM) mutR	0.5
DNA (example) extracted w/ "Qiagen DNeasy columns or other similar silica based kits"	1.5
TOTAL VOLUME OF REACTION:	15.00 μL

Comments on protocol:

- Protocol may work with other DNA extraction methods.

Strategy:

Steps	Temp (°C)	Time (m:ss)	# of Cycles
1. Initiation/Melting HOT START? ☐	94	2:00	1x
2. Denaturation	94	0:10	
3. Annealing steps 2-3-4 cycle in sequence	65 (↓1°C/cycle)	0:30	10x
4. Elongation	68	2:00	
5. Denaturation	94	0:15	
6. Annealing steps 5-6-7 cycle in sequence	55	0:30	25x
7. Elongation	68	2:00 (↑20sec/cycle)	
8. Finish	4	∞	n/a

Primers:

Name	Nucleotide Sequence (5' - 3')	Agarose: 1.5% V: 90
1. CR_Bnpl_comF	CCACTACCATCAAGTGGATTAGC	Estimated Running Time: 90 min.
2. CR_Bnpl_wtR*	GAGCAAAGGATGACAGAGAGG	Primer Combination Band (bp) Genotype
3. CR_Bnpl_mutR	ATCTCTGTGGATTCAAAGTCAGC	1 & 2,1 & 3 389, 997 wildtype
		1 & 3 538 mutant

Electrophoresis Protocol:

Allele Description: Exon 2 ENSMUSE00000176365 and flanking splicing regions were constitutively deleted from the Bnpl Gene ENSMUSG00000028115 using CRISPR Cas9 gene editing technology in mouse zygotes. The subsequent 459bp deletion from Chr 3: 95156997 - 95157455 GRCm39 was screened by PCR analysis. The selected founder animal was backcrossed to C57BL6/N to produce sequence confirmed heterozygous animals for establishing and archiving the line.

*wtR primer untested (ePCR verified)

