

GENOTYPING PROTOCOL

MUTANT MOUSE RESOURCE & RESEARCH CENTER: UC DAVIS

mmrrc@ucdavis.edu

530-754-MMRRC

Protocol Name: CR11537 Actmap (BC024978) EXDEL

Protocol: GoTaq® G2 Colorless Master Mix(Promega)

Reagent/Constituent	Volume (µL)
Water	4.5
GoTaq® G2 Colorless Master Mix,2X	7.5
Primer 1. (stock concentration is 20µM) comF	0.5
Primer 2. (stock concentration is 20µM) wtR	0.5
Primer 3. (stock concentration is 20µM) mutR	0.5
DNA (example) extracted w/ "Qiagen DNeasy columns or other similar silica based kits"	1.5
TOTAL VOLUME OF REACTION:	15.00 µL

Comments on protocol:

- Protocol may work with other DNA extraction methods.

Strategy:

Steps	Temp (°C)	Time (m:ss)	# of Cycles
1. Initiation/Melting HOT START? ☐	94	2:00	1x
2. Denaturation	94	0:10	
3. Annealing steps 2-3-4 cycle in sequence	65 (↓1°C/cycle)	0:30	10x
4. Elongation	68	2:00	
5. Denaturation	94	0:15	
6. Annealing steps 5-6-7 cycle in sequence	55	0:30	25x
7. Elongation	68	2:00 (↑20sec/cycle)	
8. Finish	4	∞	n/a

Primers:

Name	Nucleotide Sequence (5' - 3')	Agarose: 1.5% V: 90
1. CR_BC024978_comF	CCTCAGTAAGAACCCTTCGCTTGG	Estimated Running Time: 90 min.
2. CR_BC024978_wtR*	GGCTGGTAGGTGGGAGTTAAACC	Primer Combination Band (bp) Genotype
3. CR_BC024978_mutR	GCCTCTGGGGAAAAGAAGTCC	1 & 2, 1 & 3 298, 858 wildtype
		1 & 3 340 mutant

Electrophoresis Protocol:

Allele Description: Exon 4 ENSMUSE00000676543 and flanking splicing regions were constitutively deleted from the BC024978 Gene ENSMUSG00000078786 using CRISPR Cas9 gene editing technology in mouse zygotes. The subsequent 518bp deletion from Chr 7: 26900251 - 26900768 GRCm39 was screened by PCR analysis. The selected founder animal was backcrossed to C57BL6/N to produce sequence confirmed heterozygous animals for establishing and archiving the line.

*wtR primer untested (ePCR verified)

