

GENOTYPING PROTOCOL

MUTANT MOUSE RESOURCE & RESEARCH CENTER: UC DAVIS

mmrrc@ucdavis.edu

530-754-MMRRC

Protocol Name: CR11557 Fbxw11 EXDEL

Protocol: GoTaq® G2 Colorless Master Mix(Promega)

Reagent/Constituent	Volume (μL)
Water	4.5
GoTaq® G2 Colorless Master Mix,2X	7.5
Primer 1. (stock concentration is 20μM) comF	0.5
Primer 2. (stock concentration is 20μM) wtR	0.5
Primer 3. (stock concentration is 20μM) mutR	0.5
DNA (example) extracted w/ "Qiagen DNeasy columns or other similar silica based kits"	1.5
TOTAL VOLUME OF REACTION:	15.00 μL

Comments on protocol:

- Protocol may work with other DNA extraction methods.

Strategy:

Steps	Temp (°C)	Time (m:ss)	# of Cycles
1. Initiation/Melting HOT START? ☐	94	2:00	1x
2. Denaturation	94	0:10	
3. Annealing steps 2-3-4 cycle in sequence	65 (↓1°C/cycle)	0:30	10x
4. Elongation	68	2:00	
5. Denaturation	94	0:15	
6. Annealing steps 5-6-7 cycle in sequence	55	0:30	25x
7. Elongation	68	2:00 (↑20sec/cycle)	
8. Finish	4	∞	n/a

Primers:

Name	Nucleotide Sequence (5' - 3')	Agarose: 1.5%	V: 90
1. CR_Fbxw11_comF	GAGAGACATTAACCTTTAATTTTGAGG	Estimated Running Time: 90 min.	
2. CR_Fbxw11_wtR*	CAAATTAAGCTCTTTTGTTTCATCC	Primer Combination	Band (bp)
3. CR_Fbxw11_mutR	CATGTAAAATACAGATTGTCTACATACC	1 & 2, 1 & 3	290, 965
		1 & 3	308
			Genotype
			wildtype
			mutant

Allele Description: Exon 4 ENSMUSE00001223797 and flanking splicing regions were constitutively deleted from the Fbxw11 Gene ENSMUSG00000020271 using CRISPR Cas9 gene editing technology in mouse zygotes. The subsequent 657bp deletion from Chr 11: 32661512 - 32662168 GRCm39 was screened by PCR analysis. The selected founder animal was backcrossed to C57BL6/N to produce sequence confirmed heterozygous animals for establishing and archiving the line.

*wtR primer untested (ePCR verified)

