

GENOTYPING PROTOCOL

MUTANT MOUSE RESOURCE & RESEARCH CENTER: UC DAVIS

mmrrc@ucdavis.edu

530-754-MMRRC

Protocol Name: CR11560 Mfsd13a EXDEL

Protocol: GoTaq® G2 Colorless Master Mix(Promega)

Reagent/Constituent	Volume (µL)
Water	4.5
GoTaq® G2 Colorless Master Mix,2X	7.5
Primer 1. (stock concentration is 20µM) comF	0.5
Primer 2. (stock concentration is 20µM) wtR	0.5
Primer 3. (stock concentration is 20µM) mutR	0.5
DNA (example) extracted w/ "Qiagen DNeasy columns or other similar silica based kits"	1.5
TOTAL VOLUME OF REACTION:	15.00 µL

Comments on protocol:

- Protocol may work with other DNA extraction methods.

Strategy:

Steps	Temp (°C)	Time (m:ss)	# of Cycles
1. Initiation/Melting HOT START? ☐	94	2:00	1x
2. Denaturation	94	0:10	
3. Annealing steps 2-3-4 cycle in sequence	65 (↓1°C/cycle)	0:30	10x
4. Elongation	68	2:00	
5. Denaturation	94	0:15	
6. Annealing steps 5-6-7 cycle in sequence	55	0:30	25x
7. Elongation	68	2:00 (↑20sec/cycle)	
8. Finish	4	∞	n/a

Primers:

Name	Nucleotide Sequence (5' - 3')	Agarose: 1.5%	V: 90
1. CR_Mfsd13a_comF	TCCAGTGTAGGCTTTTGG	Estimated Running Time: 90 min.	
2. CR_Mfsd13a_wtR*	CTTTGTGCATGTTCTCTGC	Primer Combination	Band (bp)
3. CR_Mfsd13a_mutR	TACACCCCAAGTCTCCAACC	1 & 2, 1 & 3	484,992
		1 & 3	589
			mutant

Electrophoresis Protocol:

Allele Description: Exon 3 ENSMUSE00001279320 and flanking splicing regions were constitutively deleted from the Mfsd13a Gene ENSMUSG00000025227 using CRISPR Cas9 gene editing technology in mouse zygotes. The subsequent 403bp deletion from Chr 19: 46355482 - 46355884 GRCm39 was screened by PCR analysis. The selected founder animal was backcrossed to C57BL6/N to produce sequence confirmed heterozygous animals for establishing and archiving the line.

*wtR primer untested (ePCR verified)

