

GENOTYPING PROTOCOL

MUTANT MOUSE RESOURCE & RESEARCH CENTER: UC DAVIS

mmrrc@ucdavis.edu

530-754-MMRRC

Protocol Name: CR11558 Kdm2a EXDEL

Protocol: GoTaq® G2 Colorless Master Mix(Promega)

Reagent/Constituent	Volume (μL)
Water	4.5
GoTaq® G2 Colorless Master Mix,2X	7.5
Primer 1. (stock concentration is 20μM) comF	0.5
Primer 2. (stock concentration is 20μM) wtR	0.5
Primer 3. (stock concentration is 20μM) mutR	0.5
DNA (example) extracted w/ "Qiagen DNeasy columns or other similar silica based kits"	1.5
TOTAL VOLUME OF REACTION:	15.00 μL

Comments on protocol:

- Protocol may work with other DNA extraction methods.

Strategy:

Steps	Temp (°C)	Time (m:ss)	# of Cycles
1. Initiation/Melting HOT START? ☐	94	2:00	1x
2. Denaturation	94	0:10	
3. Annealing steps 2-3-4 cycle in sequence	65 (↓1°C/cycle)	0:30	10x
4. Elongation	68	2:00	
5. Denaturation	94	0:15	
6. Annealing steps 5-6-7 cycle in sequence	55	0:30	25x
7. Elongation	68	2:00 (↑20sec/cycle)	
8. Finish	4	∞	n/a

Primers:

Name	Nucleotide Sequence (5' - 3')	Agarose: 1.5%	V: 90
1. CR_Kdm2a_comF	GAGCTGTCAAACATGGGTACTGG	Estimated Running Time: 90 min.	
2. CR_Kdm2a_wtR*	GCTGCTAGTCCAAAGCATTACG	Primer Combination	Band (bp)
3. CR_Kdm2a_mutR	CAGAGAGCACTTACTTTATTCCAAGTCC	1 & 2, 1 & 3	432, 1488
		1 & 3	672
			wildtype
			mutant

Electrophoresis Protocol:

Allele Description: Exon 3 ENSMUSE00001246260 and flanking splicing regions were constitutively deleted from the Kdm2a Gene ENSMUSG00000054611 using CRISPR Cas9 gene editing technology in mouse zygotes. The subsequent 816bp deletion from Chr 19: 4412342 - 4413157 GRCm39 was screened by PCR analysis. The selected founder animal was backcrossed to C57BL6/N to produce sequence confirmed heterozygous animals for establishing and archiving the line.

*wtR primer untested (ePCR verified)

