GENOTYPING PROTOCOL

MUTANT MOUSE RESOURCE & RESEARCH CENTER: UC DAVIS

mmrrc@ucdavis.edu
530-754-MMRRC

Protocol Name: **CR11551 Cntnap4 EXDEL**

Protocol: ***GoTaq® G2 Colorless Master Mix(Promega)***

Reagent/Constituent	Volume (µL)
Water	4.5
GoTaq® G2 Colorless Master Mix,2X	7.5
Primer 1. (stock concentration is 20µM) comF	0.5
Primer 2. (stock concentration is 20µM) wtR	0.5
Primer 3. (stock concentration is 20µM) mutR	0.5
DNA (example) extracted w/ "Qiagen DNeasy columns or other similar silica based kits"	1.5
TOTAL VOLUME OF REACTION:	**15.00 µL**

Comments on protocol:

- Protocol may work with other DNA extraction methods.

Strategy:

Steps		Temp (ºC)	Time (m:ss)	# of Cycles
1. Initiation/Melting	HOT START? ☐	94	2:00	**1x**
2. Denaturation		94	0:10	
3. Annealing	steps 2-3-4 cycle in sequence	65 (↓1ºC/cycle)	0:30	**10x**
4. Elongation		68	2:00	
5. Denaturation		94	0:15	
6. Annealing	steps 5-6-7 cycle in sequence	55	0:30	**25x**
7. Elongation		68	2:00 (↑20sec/cycle)	
8. Finish		4	∞	n/a

Primers:

Name	Nucleotide Sequence (5' - 3')
1. CR_Cntnap4_comF	CCCTTACCAAACCCAATTCATTCC
2. CR_Cntnap4_wtR*	GTGTGCCACTTGTGTTCATTTAGAGG
3. CR_Cntnap4_mutR	CTTCAGACATAGCTCTCTACATTACAGC

Electrophoresis Protocol:

Agarose: **1.5%** **V:** **90**

Estimated Running Time: **90** min.

Primer Combination	**Band** (bp)	**Genotype**
1 & 2,1 & 3	465, 1231	wildtype
1 & 3	404	mutant

Allele Description: Exon 3 ENSMUSE00001238256 and flanking splicing regions were constitutively deleted from the Cntnap4 Gene ENSMUSG00000031772 using CRISPR Cas9 gene editing technology in mouse zygotes. The subsequent 827bp deletion from Chr 8: 113391425 - 113392251 GRCm39 was screened by PCR analysis. The selected founder animal was backcrossed to C57BL6/N to produce sequence confirmed heterozygous animals for establishing and archiving the line.

*wtR primer untested (ePCR verified)