

GENOTYPING PROTOCOL

MUTANT MOUSE RESOURCE & RESEARCH CENTER: UC DAVIS

mmrrc@ucdavis.edu

530-754-MMRRC

Protocol Name: CR11563 Mospd3 EXDEL

Protocol: GoTaq® G2 Colorless Master Mix(Promega)

Reagent/Constituent	Volume (μL)
Water	4.5
GoTaq® G2 Colorless Master Mix,2X	7.5
Primer 1. (stock concentration is 20μM) comF	0.5
Primer 2. (stock concentration is 20μM) wtR	0.5
Primer 3. (stock concentration is 20μM) mutR	0.5
DNA (example) extracted w/ "Qiagen DNeasy columns or other similar silica based kits"	1.5
TOTAL VOLUME OF REACTION:	15.00 μL

Comments on protocol:

- Protocol may work with other DNA extraction methods.

Strategy:

Steps	Temp (°C)	Time (m:ss)	# of Cycles
1. Initiation/Melting HOT START? ☐	94	2:00	1x
2. Denaturation	94	0:10	
3. Annealing steps 2-3-4 cycle in sequence	65 (↓1°C/cycle)	0:30	10x
4. Elongation	68	2:00	
5. Denaturation	94	0:15	
6. Annealing steps 5-6-7 cycle in sequence	55	0:30	25x
7. Elongation	68	2:00 (↑20sec/cycle)	
8. Finish	4	∞	n/a

Primers:

Name	Nucleotide Sequence (5' - 3')	Agarose: 1.5% V: 90
1. CR_Mospd3_comF	TCTCAGACAGAGAAGGAAGTCTCAGG	Estimated Running Time: 90 min.
2. CR_Mospd3_wtR*	ATTGTGTCCCTGTGCCCTTTCC	Primer Combination Band (bp) Genoty
3. CR_Mospd3_mutR	GCATACATACATACATACATACTCACG	1 & 2,1 & 3 326,4702 wildtype
		1 & 3 352 mutant

Electrophoresis Protocol:

Allele Description: Exon 1-6 (ENSMUSE00000387892, ENSMUSE00000343999, ENSMUSE00001240860, ENSMUSE00001216743, ENSMUSE00000306973, ENSMUSE00000379307)were constitutively deleted from the 5'UTR through the 3'UTR from the Mospd3 Gene ENSMUSG00000037221 using CRISPR Cas9 gene editing technology in mouse zygotes. The subsequent 4350bp deletion from Chr 5: 137594992 - 137599341 GRCh39 was screened by PCR analysis. The selected founder animal was backcrossed to C57BL6/N to produce sequence confirmed heterozygous animals for establishing and archiving the line.

*wtR primer untested (ePCR verified)

