

GENOTYPING PROTOCOL

MUTANT MOUSE RESOURCE & RESEARCH CENTER: UC DAVIS

mmrrc@ucdavis.edu

530-754-MMRRC

Protocol Name: CR11606 Lactb2 EXDEL

Protocol: GoTaq® G2 Colorless Master Mix(Promega)

Reagent/Constituent	Volume (µL)
Water	4.5
GoTaq® G2 Colorless Master Mix,2X	7.5
Primer 1. (stock concentration is 20µM) comF	0.5
Primer 2. (stock concentration is 20µM) wtR	0.5
Primer 3. (stock concentration is 20µM) mutR	0.5
DNA (example) extracted w/ "Qiagen DNeasy columns or other similar silica based kits"	1.5
TOTAL VOLUME OF REACTION:	15.00 µL

Comments on protocol:

- Protocol may work with other DNA extraction methods.

Strategy:

Steps	Temp (°C)	Time (m:ss)	# of Cycles
1. Initiation/Melting HOT START? <input type="checkbox"/>	94	2:00	1x
2. Denaturation	94	0:10	
3. Annealing steps 2-3-4 cycle in sequence	65 (↓1°C/cycle)	0:30	10x
4. Elongation	68	2:00	
5. Denaturation	94	0:15	
6. Annealing steps 5-6-7 cycle in sequence	55	0:30	25x
7. Elongation	68	2:00 (↑20sec/cycle)	
8. Finish	4	∞	n/a

Primers:

Name	Nucleotide Sequence (5' - 3')
1. CR_Lactb2_comF	CTGACACTGCTAAGTGGTAAAGC
2. CR_Lactb2_wtR*	CAGAGTGTCTTCTCCATGC
3. CR_Lactb2_mutR	TCTCAAGTCTGTATTTCTCAATAGTGG

Electrophoresis Protocol:

Agarose: 1.5%	V: 90
Estimated Running Time: 90 min.	
Primer Combination	Band (bp)
1 & 2, 1 & 3	295,846
1 & 3	321
	Genotype
	wildtype
	mutant

Allele Description: Exon 3 ENSMUSE00000154321 and flanking splicing regions were constitutively deleted from the Lactb2 Gene ENSMUSG00000025937 using CRISPR Cas9 gene editing technology in mouse zygotes. The subsequent 525bp deletion from Chr 1:13717537 - 13718061 GRCm39 was screened by PCR analysis. The selected founder animal was backcrossed to C57BL6/N to produce sequence confirmed heterozygous animals for establishing and archiving the line.

*wtR primer untested (ePCR verified)

