

GENOTYPING PROTOCOL

MUTANT MOUSE RESOURCE & RESEARCH CENTER: UC DAVIS

mmrrc@ucdavis.edu

530-754-MMRRC

Protocol Name: CR11678 Poldip2 EXDEL

Stock # 75984

Protocol: GoTaq® G2 Colorless Master Mix(Promega)

Reagent/Constituent	Volume (µL)
Water	4.5
GoTaq® G2 Colorless Master Mix,2X	7.5
Primer 1. (stock concentration is 20µM) comF	0.5
Primer 2. (stock concentration is 20µM) wtR	0.5
Primer 3. (stock concentration is 20µM) mutR	0.5
DNA (example) extracted w/ "Qiagen DNeasy columns or other similar silica based kits"	1.5
TOTAL VOLUME OF REACTION:	15.00 µL

Comments on protocol:

- Protocol may work with other DNA extraction methods.

Strategy:

Steps	Temp (°C)	Time (m:ss)	# of Cycles
1. Initiation/Melting HOT START? ☐	94	2:00	1x
2. Denaturation	94	0:10	
3. Annealing steps 2-3-4 cycle in sequence	65 (↓1°C/cycle)	0:30	10x
4. Elongation	68	2:00	
5. Denaturation	94	0:15	
6. Annealing steps 5-6-7 cycle in sequence	55	0:30	25x
7. Elongation	68	2:00 (↑20sec/cycle)	
8. Finish	4	∞	n/a

Primers:

Name	Nucleotide Sequence (5' - 3')	Agarose: 1.5%	V: 90
1. CR_Poldip2_comF	ACACACTAGGCATGTCTCTCAGG	Estimated Running Time: 90 min.	
2. CR_Poldip2_wtR*	AGTGAGACCCTGAGAAGACAGC	Primer Combination	Band (bp)
3. CR_Poldip2_mutR	CGTGGGTTACAGAATGAGTCAAGG	1 & 2, 1 & 3	396, 774
		1 & 3	359
			mutant

Electrophoresis Protocol:

Allele Description: Exon 2 ENSMUSE00000109815 and flanking splicing regions were constitutively deleted from the Poldip2 Gene ENSMUSG00000001100 using CRISPR Cas9 gene editing technology in mouse zygotes. The subsequent 415bp deletion from Chr11:78404540-78404954 GRCm39 was screened by PCR analysis. The selected founder animal was backcrossed to C57BL6/N to produce sequence confirmed heterozygous animals for establishing and archiving the line.

*wtR primer untested (ePCR verified)

