

GENOTYPING PROTOCOL

MUTANT MOUSE RESOURCE & RESEARCH CENTER: UC DAVIS

mmrrc@ucdavis.edu

530-754-MMRRC

Protocol Name: CR11686 Tyw5 EXDEL

Stock # 75991

Protocol: GoTaq® G2 Colorless Master Mix(Promega)

Reagent/Constituent	Volume (μL)
Water	4.5
GoTaq® G2 Colorless Master Mix,2X	7.5
Primer 1. (stock concentration is 20μM) comF	0.5
Primer 2. (stock concentration is 20μM) wtR	0.5
Primer 3. (stock concentration is 20μM) mutR	0.5
DNA (example) extracted w/ "Qiagen DNeasy columns or other similar silica based kits"	1.5
TOTAL VOLUME OF REACTION:	15.00 μL

Comments on protocol:

- Protocol may work with other DNA extraction methods.

Strategy:

Steps	Temp (°C)	Time (m:ss)	# of Cycles
1. Initiation/Melting HOT START? ☐	94	2:00	1x
2. Denaturation	94	0:10	
3. Annealing steps 2-3-4 cycle in sequence	65 (↓1°C/cycle)	0:30	10x
4. Elongation	68	2:00	
5. Denaturation	94	0:15	
6. Annealing steps 5-6-7 cycle in sequence	55	0:30	25x
7. Elongation	68	2:00 (↑20sec/cycle)	
8. Finish	4	∞	n/a

Primers:

Electrophoresis Protocol:

Name	Nucleotide Sequence (5' - 3')	Agarose: 1.5%	V: 90
1. CR_Tyw5_comF	GTAGTTTTAGTAACATATCTGGGGCAG	Estimated Running Time: 90 min.	
2. CR_Tyw5_wtR*	AACAACCTTCCTTGGTATCAGAGTGTGG	Primer Combination	Band (bp)
3. CR_Tyw5_mutR	GCAACTACCACACAGGATTGGAAAGTAG	1 & 2, 1 & 3	519, 1483
		1 & 3	566
			mutant

Allele Description: Exon 2 ENSMUSE00000972411 and flanking splicing regions were constitutively deleted from the Tyw5 Gene ENSMUSG00000048495 using CRISPR Cas9 gene editing technology in mouse zygotes. The subsequent 917bp deletion from Chr 1: 57440050 – 57440966 GRCh39 was screened by PCR analysis. The selected founder animal was backcrossed to C57BL6/N to produce sequence confirmed heterozygous animals for establishing and archiving the line.

*wtR primer untested (ePCR verified)

