

GENOTYPING PROTOCOL

MUTANT MOUSE RESOURCE & RESEARCH CENTER: UC DAVIS

mmrrc@ucdavis.edu

530-754-MMRRC

Protocol Name: CR11687 Vstm2l EXDEL

Protocol: GoTaq® G2 Colorless Master Mix(Promega)

Reagent/Constituent	Volume (μL)
Water	4.5
GoTaq® G2 Colorless Master Mix,2X	7.5
Primer 1. (stock concentration is 20μM) comF	0.5
Primer 2. (stock concentration is 20μM) wtR	0.5
Primer 3. (stock concentration is 20μM) mutR	0.5
DNA (example) extracted w/ "Qiagen DNeasy columns or other similar silica based kits"	1.5
TOTAL VOLUME OF REACTION:	15.00 μL

Comments on protocol:

- Protocol may work with other DNA extraction methods.

Strategy:

Steps	Temp (°C)	Time (m:ss)	# of Cycles
1. Initiation/Melting HOT START? ☐	94	2:00	1x
2. Denaturation	94	0:10	
3. Annealing steps 2-3-4 cycle in sequence	65 (↓1°C/cycle)	0:30	10x
4. Elongation	68	2:00	
5. Denaturation	94	0:15	
6. Annealing steps 5-6-7 cycle in sequence	55	0:30	25x
7. Elongation	68	2:00 (↑20sec/cycle)	
8. Finish	4	∞	n/a

Primers:

Name	Nucleotide Sequence (5' - 3')	Agarose: 1.5%	V: 90
1. CR_Vstm2l_comF	GAACGGGTTGTTCTGTGACTTGG	Estimated Running Time: 90 min.	
2. CR_Vstm2l_wtR*	GAAGAGTGCTGTGTGTGGAAAGG	Primer Combination	Band (bp)
3. CR_Vstm2l_mutR	GGAACTCAGGTCTTATCAATGACTCAGC	1 & 2, 1 & 3	585,1318
		1 & 3	702
			mutant

Electrophoresis Protocol:

Allele Description: Exon 2 ENSMUSE00000316305 and flanking splicing regions were constitutively deleted from the Vstm2l Gene ENSMUSG00000037843 using CRISPR Cas9 gene editing technology in mouse zygotes. The subsequent 616bp deletion from Chr2: 157776969 - 157777584 GRCm39 was screened by PCR analysis. The selected founder animal was backcrossed to C57BL6/N to produce sequence confirmed heterozygous animals for establishing and archiving the line.

*wtR primer untested (ePCR verified)

