GENOTYPING PROTOCOL

MUTANT MOUSE RESOURCE & RESEARCH CENTER: UC DAVIS

mmrrc@ucdavis.edu
530-754-MMRRC

Protocol Name: **CR11688 Wbp4 EXDEL**

Protocol: ***GoTaq® G2 Colorless Master Mix(Promega)***

Reagent/Constituent	Volume (µL)
Water	4.5
GoTaq® G2 Colorless Master Mix,2X	7.5
Primer 1. (stock concentration is 20µM) comF	0.5
Primer 2. (stock concentration is 20µM) wtR	0.5
Primer 3. (stock concentration is 20µM) mutR	0.5
DNA (example) extracted w/ "Qiagen DNeasy columns or other similar silica based kits"	1.5
TOTAL VOLUME OF REACTION:	**15.00 µL**

Comments on protocol:

- Protocol may work with other DNA extraction methods.

Strategy:

Steps		Temp (ºC)	Time (m:ss)	# of Cycles
1. Initiation/Melting	HOT START? ☐	94	2:00	**1x**
2. Denaturation		94	0:10	
3. Annealing	steps 2-3-4 cycle in sequence	65 (↓1ºC/cycle)	0:30	**10x**
4. Elongation		68	2:00	
5. Denaturation		94	0:15	
6. Annealing	steps 5-6-7 cycle in sequence	55	0:30	**25x**
7. Elongation		68	2:00 (↑20sec/cycle)	
8. Finish		4	∞	n/a

Primers:

Name	Nucleotide Sequence (5' - 3')
1. CR_Wbp4_comF	GGTGGTTGTTTTAGTTTGGGGAGATAGG
2. CR_Wbp4_wtR*	CCTTTTCTGGAGTATCTGAGGACAGC
3. CR_Wbp4_mutR	TTACCTAAACTCTCAATAAACTGCCTGC

Electrophoresis Protocol:

Agarose: **1.5%** **V:** **90**

Estimated Running Time: **90** min.

Primer Combination	**Band** (bp)	**Genotype**
1 & 2,1 & 3	467,2230	wildtype
1 & 3	515	mutant

Allele Description: Exon 3-4 (ENSMUSE00000123034, ENSMUSE00000123033) and flanking splicing regions were constitutively deleted from the Wbp4 Gene ENSMUSG00000022023 using CRISPR Cas9 gene editing technology in mouse zygotes. The subsequent 1715bp deletion from Chr 14: 79713232 - 79714946 GRCm39 was screened by PCR analysis. The selected founder animal was backcrossed to C57BL6/N to produce sequence confirmed heterozygous animals for establishing and archiving the line.

*wtR primer untested (ePCR verified)