

GENOTYPING PROTOCOL

MUTANT MOUSE RESOURCE & RESEARCH CENTER: UC DAVIS

mmrrc@ucdavis.edu

530-754-MMRRC

Protocol Name: CR11702 Cda EXDEL

Protocol: GoTaq® G2 Colorless Master Mix(Promega)

Reagent/Constituent	Volume (µL)
Water	4.5
GoTaq® G2 Colorless Master Mix,2X	7.5
Primer 1. (stock concentration is 20µM) comF	0.5
Primer 2. (stock concentration is 20µM) wtR	0.5
Primer 3. (stock concentration is 20µM) mutR	0.5
DNA (example) extracted w/ "Qiagen DNeasy columns or other similar silica based kits"	1.5
TOTAL VOLUME OF REACTION:	15.00 µL

Comments on protocol:

- Protocol may work with other DNA extraction methods.

Strategy:

Steps	Temp (°C)	Time (m:ss)	# of Cycles
1. Initiation/Melting HOT START? <input type="checkbox"/>	94	2:00	1x
2. Denaturation	94	0:10	
3. Annealing steps 2-3-4 cycle in sequence	65 (↓1°C/cycle)	0:30	10x
4. Elongation	68	2:00	
5. Denaturation	94	0:15	
6. Annealing steps 5-6-7 cycle in sequence	55	0:30	25x
7. Elongation	68	2:00 (↑20sec/cycle)	
8. Finish	4	∞	n/a

Primers:

Electrophoresis Protocol:

Name	Nucleotide Sequence (5' - 3')	Agarose: 1.5%	V: 90
1. CR_Cda_comF	GTGGTGGGAGTGCTCTCATTAGG	Estimated Running Time: 90 min.	
2. CR_Cda_wtR*	GAAACCTCAGTTGAGAGAATGTGTGC	Primer Combination	Band (bp)
3. CR_Cda_mutR	CCTCAAGAGCAAGTGTCAAGTGC	1 & 2, 1 & 3	416, 1113
		1 & 3	479
			Genotype
			wildtype
			mutant

Allele Description: Exon 2 ENSMUSE00000181843 and flanking splicing regions were constitutively deleted from the Cda Gene ENSMUSG00000028755 using CRISPR Cas9 gene editing technology in mouse zygotes. The subsequent 634bp deletion from Chr 4: 138078255 - 138078888 GRCm39 was screened by PCR analysis. The selected founder animal was backcrossed to C57BL6/N to produce sequence confirmed heterozygous animals for establishing and archiving the line.

*wtR primer untested (ePCR verified)

