

GENOTYPING PROTOCOL

MUTANT MOUSE RESOURCE & RESEARCH CENTER: UC DAVIS

mmrrc@ucdavis.edu

530-754-MMRRC

Protocol Name: CR11726 Smim13 EXDEL

Protocol: GoTaq® G2 Colorless Master Mix(Promega)

Reagent/Constituent	Volume (µL)
Water	4.5
GoTaq® G2 Colorless Master Mix,2X	7.5
Primer 1. (stock concentration is 20µM) comF	0.5
Primer 2. (stock concentration is 20µM) wtR	0.5
Primer 3. (stock concentration is 20µM) mutR	0.5
DNA (example) extracted w/ "Qiagen DNeasy columns or other similar silica based kits"	1.5
TOTAL VOLUME OF REACTION:	15.00 µL

Comments on protocol:

- Protocol may work with other DNA extraction methods.

Strategy:

Steps	Temp (°C)	Time (m:ss)	# of Cycles
1. Initiation/Melting HOT START? <input type="checkbox"/>	94	2:00	1x
2. Denaturation	94	0:10	
3. Annealing steps 2-3-4 cycle in sequence	65 (↓1°C/cycle)	0:30	10x
4. Elongation	68	2:00	
5. Denaturation	94	0:15	
6. Annealing steps 5-6-7 cycle in sequence	55	0:30	25x
7. Elongation	68	2:00 (↑20sec/cycle)	
8. Finish	4	∞	n/a

Primers:

Electrophoresis Protocol:

Name	Nucleotide Sequence (5' - 3')	Agarose: 1.5%	V: 90
1. CR_Smim13_comF	GAACAAGAATCAGCAGTCTCTAAGTTGG	Estimated Running Time: 90 min.	
2. CR_Smim13_wtR*	CGGGGAAAGAGTGATCTGGAGG	Primer Combination	Band (bp)
3. CR_Smim13_mutR	GCATTATGGCAACCAAGGACAAACC	1 & 2, 1 & 3	379,997
		1 & 3	459
			wildtype
			mutant

Allele Description: Exon 1 ENSMUSE00000915565 and flanking splicing regions were constitutively deleted from the Smim13 Gene ENSMUSG0000091264 using CRISPR Cas9 gene editing technology in mouse zygotes. The subsequent 538bp deletion from Chr 13: 41403276- 41403813 GRcm39 was sequence verified. The selected founder animal was backcrossed to C57BL6/N to produce sequence confirmed heterozygous animals for establishing and archiving the line.

*wtR primer untested (ePCR verified)

