

GENOTYPING PROTOCOL

MUTANT MOUSE RESOURCE & RESEARCH CENTER: UC DAVIS

mmrrc@ucdavis.edu

530-754-MMRRC

Protocol Name: CR11737 Ttc3 EXDEL

Protocol: GoTaq® G2 Colorless Master Mix(Promega)

Reagent/Constituent	Volume (μL)
Water	4.5
GoTaq® G2 Colorless Master Mix,2X	7.5
Primer 1. (stock concentration is 20μM) comF	0.5
Primer 2. (stock concentration is 20μM) wtR	0.5
Primer 3. (stock concentration is 20μM) mutR	0.5
DNA (example) extracted w/ "Qiagen DNeasy columns or other similar silica based kits"	1.5
TOTAL VOLUME OF REACTION:	15.00 μL

Comments on protocol:

- Protocol may work with other DNA extraction methods.

Strategy:

Steps	Temp (°C)	Time (m:ss)	# of Cycles
1. Initiation/Melting HOT START? ☐	94	2:00	1x
2. Denaturation	94	0:10	
3. Annealing steps 2-3-4 cycle in sequence	65 (↓1°C/cycle)	0:30	10x
4. Elongation	68	2:00	
5. Denaturation	94	0:15	
6. Annealing steps 5-6-7 cycle in sequence	55	0:30	25x
7. Elongation	68	2:00 (↑20sec/cycle)	
8. Finish	4	∞	n/a

Primers:

Name	Nucleotide Sequence (5' - 3')	Electrophoresis Protocol:		
1. CR_Ttc3_comF	CAGTCTGTTGTGTGAAAGGCTGG	Agarose: 1.5%	V: 90	
2. CR_Ttc3_wtR*	GCCAGGTGTTCTTCAAAACAATGG	Estimated Running Time: 90 min.		
3. CR_Ttc3_mutR	CTGTGACAAAGTGAGCTTTCTCTGC	Primer Combination	Band (bp)	Genotype
		1 & 2, 1 & 3	624,1087	wildtype
		1 & 3	601	mutant

Allele Description: Exon 11 ENSMUSE00001260102 and flanking splicing regions were constitutively deleted from the Ttc3 Gene ENSMUSG00000040785 using CRISPR Cas9 gene editing technology in mouse zygotes. The subsequent 486bp deletion from Chr 16: 94203963-94204448 GRCm39 was sequence verified. The selected founder animal was backcrossed to C57BL6/N to produce sequence confirmed heterozygous animals for establishing and archiving the line.

*wtR primer untested (ePCR verified)

