

GENOTYPING PROTOCOL

MUTANT MOUSE RESOURCE & RESEARCH CENTER: UC DAVIS

mmrrc@ucdavis.edu

530-754-MMRRC

Protocol Name: CR11751 Aunip EXDEL

Protocol: GoTaq® G2 Colorless Master Mix(Promega)

Reagent/Constituent	Volume (μL)
Water	4.5
GoTaq® G2 Colorless Master Mix,2X	7.5
Primer 1. (stock concentration is 20μM) comF	0.5
Primer 2. (stock concentration is 20μM) wtR	0.5
Primer 3. (stock concentration is 20μM) mutR	0.5
DNA (example) extracted w/ "Qiagen DNeasy columns or other similar silica based kits"	1.5
TOTAL VOLUME OF REACTION:	15.00 μL

Comments on protocol:

- Protocol may work with other DNA extraction methods.

Strategy:

Steps	Temp (°C)	Time (m:ss)	# of Cycles
1. Initiation/Melting HOT START? ☐	94	2:00	1x
2. Denaturation	94	0:10	
3. Annealing steps 2-3-4 cycle in sequence	65 (↓1°C/cycle)	0:30	10x
4. Elongation	68	2:00	
5. Denaturation	94	0:15	
6. Annealing steps 5-6-7 cycle in sequence	55	0:30	25x
7. Elongation	68	2:00 (↑20sec/cycle)	
8. Finish	4	∞	n/a

Primers:

Name	Nucleotide Sequence (5' - 3')	Agarose: 1.5%	V: 90
1. CR_Aunip_comF	GGTTAGGAGCAGAGAAAGTAGACACC	Estimated Running Time: 90 min.	
2. CR_Aunip_wtR*	CTGTAACAGCAGTGATTCTTGAAGC	Primer Combination	Band (bp)
3. CR_Aunip_mutR	CCTCAATTACGCTTTCTAGTCAGACC	1 & 2, 1 & 3	474, 1780
		1 & 3	436
			mutant

Electrophoresis Protocol:

Allele Description: Exon 2 ENSMUSE00000667837 and flanking splicing regions were constitutively deleted from the Aunip Gene ENSMUSG00000078521 using CRISPR Cas9 gene editing technology in mouse zygotes. The subsequent 1344bp deletion from Chr4:134248435-134249778 GRCm39 was sequence verified. The selected founder animal was backcrossed to C57BL6/N to produce sequence confirmed heterozygous animals for establishing and archiving the line.

*wtR primer untested (ePCR verified)

