

GENOTYPING PROTOCOL

MUTANT MOUSE RESOURCE & RESEARCH CENTER: UC DAVIS

mmrrc@ucdavis.edu

530-754-MMRRC

Protocol Name: CR11759 Dgcr6 EXDEL

Protocol: GoTaq® G2 Colorless Master Mix(Promega)

Reagent/Constituent	Volume (μL)
Water	4.5
GoTaq® G2 Colorless Master Mix,2X	7.5
Primer 1. (stock concentration is 20μM) comF	0.5
Primer 2. (stock concentration is 20μM) wtR	0.5
Primer 3. (stock concentration is 20μM) mutR	0.5
DNA (example) extracted w/ "Qiagen DNeasy columns or other similar silica based kits"	1.5
TOTAL VOLUME OF REACTION:	15.00 μL

Comments on protocol:

- Protocol may work with other DNA extraction methods.

Strategy:

Steps	Temp (°C)	Time (m:ss)	# of Cycles
1. Initiation/Melting HOT START? ☐	94	2:00	1x
2. Denaturation	94	0:10	
3. Annealing steps 2-3-4 cycle in sequence	65 (↓1°C/cycle)	0:30	10x
4. Elongation	68	2:00	
5. Denaturation	94	0:15	
6. Annealing steps 5-6-7 cycle in sequence	55	0:30	25x
7. Elongation	68	2:00 (↑20sec/cycle)	
8. Finish	4	∞	n/a

Primers:

Name	Nucleotide Sequence (5' - 3')	Agarose: 1.5%	V: 90
1. CR_Dgcr6_comF	CTGTCCCATGAATCTCCTCTAGG	Estimated Running Time: 90 min.	
2. CR_Dgcr6_wtR*	CTGTGTAGGTTAGGCAGGTATAGC	Primer Combination	Band (bp)
3. CR_Dgcr6_mutR	CAGAGATGCTTGTGGCCAGG	1 & 2, 1 & 3	551, 1017
		1 & 3	282
			wildtype
			mutant

Electrophoresis Protocol:

Allele Description: Exon 2 ENSMUSE00000367792 and flanking splicing regions were constitutively deleted from the Dgcr6 Gene ENSMUSG00000003531 using CRISPR Cas9 gene editing technology in mouse zygotes. The subsequent 733bp deletion from Chr16: 17886715- 17887447 GRCm39 was sequence verified. The selected founder animal was backcrossed to C57BL6/N to produce sequence confirmed heterozygous animals for establishing and archiving the line.

*wtR primer untested (ePCR verified)

