

GENOTYPING PROTOCOL

MUTANT MOUSE RESOURCE & RESEARCH CENTER: UC DAVIS

mmrrc@ucdavis.edu

530-754-MMRRC

Protocol Name: **CR11754 BC005624 EXDEL**

Protocol: **GoTaq® G2 Colorless Master Mix(Promega)**

Reagent/Constituent	Volume (μL)
Water	4.5
GoTaq® G2 Colorless Master Mix,2X	7.5
Primer 1. (stock concentration is 20μM) comF	0.5
Primer 2. (stock concentration is 20μM) wtR	0.5
Primer 3. (stock concentration is 20μM) mutR	0.5
DNA (example) extracted w/ "Qiagen DNeasy columns or other similar silica based kits"	1.5
TOTAL VOLUME OF REACTION:	15.00 μL

Comments on protocol:

- Protocol may work with other DNA extraction methods.

Strategy:

Steps	Temp (°C)	Time (m:ss)	# of Cycles
1. Initiation/Melting HOT START? ☐	94	2:00	1x
2. Denaturation	94	0:10	
3. Annealing steps 2-3-4 cycle in sequence	65 (↓1°C/cycle)	0:30	10x
4. Elongation	68	2:00	
5. Denaturation	94	0:15	
6. Annealing steps 5-6-7 cycle in sequence	55	0:30	25x
7. Elongation	68	2:00 (↑20sec/cycle)	
8. Finish	4	∞	n/a

Primers:

Electrophoresis Protocol:

Name	Nucleotide Sequence (5' - 3')	Agarose: 1.5%	V: 90
1. CR_BC005624_comF	GGCAGAACCTTACTCTGTGATTTCC	Estimated Running Time: 90 min.	
2. CR_BC005624_wtR*	CTCCTCTTGAACCTTCTCTCTACC	Primer Combination	Band (bp)
3. CR_BC005624_mutR	GTACCTTGGCACACTTATGGTATGC	1 & 2, 1 & 3	550, 1769
		1 & 3	861
			mutant

Allele Description: Exon 3 ENSMUSE00000163575 and flanking splicing regions were constitutively deleted from the BC005624 Gene ENSMUSG00000026851 using CRISPR Cas9 gene editing technology in mouse zygotes. The subsequent 908bp deletion from Chr2: 30869325- 30870232 GRCm39 was sequence verified. The selected founder animal was backcrossed to C57BL6/N to produce sequence confirmed heterozygous animals for establishing and archiving the line.

*wtR primer untested (ePCR verified)

