

GENOTYPING PROTOCOL

MUTANT MOUSE RESOURCE & RESEARCH CENTER: UC DAVIS

mmrrc@ucdavis.edu

530-754-MMRRC

Protocol Name: CR11756 Ccdc43 EXDEL

Protocol: GoTaq® G2 Colorless Master Mix(Promega)

Reagent/Constituent	Volume (μL)
Water	4.5
GoTaq® G2 Colorless Master Mix,2X	7.5
Primer 1. (stock concentration is 20μM) comF	0.5
Primer 2. (stock concentration is 20μM) wtR	0.5
Primer 3. (stock concentration is 20μM) mutR	0.5
DNA (example) extracted w/ "Qiagen DNeasy columns or other similar silica based kits"	1.5
TOTAL VOLUME OF REACTION:	15.00 μL

Comments on protocol:

- Protocol may work with other DNA extraction methods.

Strategy:

Steps	Temp (°C)	Time (m:ss)	# of Cycles
1. Initiation/Melting HOT START? ☐	94	2:00	1x
2. Denaturation	94	0:10	
3. Annealing steps 2-3-4 cycle in sequence	65 (↓1°C/cycle)	0:30	10x
4. Elongation	68	2:00	
5. Denaturation	94	0:15	
6. Annealing steps 5-6-7 cycle in sequence	55	0:30	25x
7. Elongation	68	2:00 (↑20sec/cycle)	
8. Finish	4	∞	n/a

Primers:

Name	Nucleotide Sequence (5' - 3')	Agarose: 1.5% V: 90
1. CR_Ccdc43_comF	CAGACAGATCTCTGTGAGTTTGAGTCC	Estimated Running Time: 90 min.
2. CR_Ccdc43_wtR*	GTGACATCCCGTGTTTCTGACC	Primer Combination Band (bp) Genotype
3. CR_Ccdc43_mutR	CAGGTGGATTGGAGAGACAAAGG	1 & 2,1 & 3 411,942 wildtype
		1 & 3 272 mutant

Electrophoresis Protocol:

Allele Description: Exon 2 ENSMUSE00001241837 and flanking splicing regions were constitutively deleted from the Ccdc43 Gene ENSMUSG00000020925 using CRISPR Cas9 gene editing technology in mouse zygotes. The subsequent 670bp deletion from Chr11: 102582560- 102583229 GRCm39 was sequence verified. The selected founder animal was backcrossed to C57BL6/N to produce sequence confirmed heterozygous animals for establishing and archiving the line.

*wtR primer untested (ePCR verified)

