

GENOTYPING PROTOCOL

MUTANT MOUSE RESOURCE & RESEARCH CENTER: UC DAVIS

mmrrc@ucdavis.edu

530-754-MMRRC

Protocol Name: CR11731 Spata31g1 EXDEL

Protocol: GoTaq® G2 Colorless Master Mix(Promega)

Reagent/Constituent	Volume (µL)
Water	4.5
GoTaq® G2 Colorless Master Mix,2X	7.5
Primer 1. (stock concentration is 20µM) comF	0.5
Primer 2. (stock concentration is 20µM) wtR	0.5
Primer 3. (stock concentration is 20µM) mutR	0.5
DNA (example) extracted w/ "Qiagen DNeasy columns or other similar silica based kits"	1.5
TOTAL VOLUME OF REACTION:	15.00 µL

Comments on protocol:

- Protocol may work with other DNA extraction methods.

Strategy:

Steps	Temp (°C)	Time (m:ss)	# of Cycles
1. Initiation/Melting HOT START? <input type="checkbox"/>	94	2:00	1x
2. Denaturation	94	0:10	
3. Annealing steps 2-3-4 cycle in sequence	65 (↓1°C/cycle)	0:30	10x
4. Elongation	68	2:00	
5. Denaturation	94	0:15	
6. Annealing steps 5-6-7 cycle in sequence	55	0:30	25x
7. Elongation	68	2:00 (↑20sec/cycle)	
8. Finish	4	∞	n/a

Primers:

Name	Nucleotide Sequence (5' - 3')	Agarose: 1.5%	V: 90
1. CR_Spata31g1_comF	AGATGCTGGTGTCTAAATGAGAACTGG	Estimated Running Time: 90 min.	
2. CR_Spata31g1_wtR*	AGGGTCTGGAAAGTAGGGAAGG	Primer Combination	Band (bp)
3. CR_Spata31g1_mutR	TCATCTTGCCAGCCCTGAAACC	1 & 2, 1 & 3	754,3998
		1 & 3	622
			wildtype
			mutant

Electrophoresis Protocol:

Allele Description: Exon 2 ENSMUSE00000605002 and flanking splicing regions were constitutively deleted from the Spata31g1 Gene ENSMUSG00000028451 using CRISPR Cas9 gene editing technology in mouse zygotes. The subsequent 3376bp deletion from Chr 4: 42970846- 42974221 GRcm39 was sequence verified. The selected founder animal was backcrossed to C57BL6/N to produce sequence confirmed heterozygous animals for establishing and archiving the line.

*wtR primer untested (ePCR verified)

