

GENOTYPING PROTOCOL

MUTANT MOUSE RESOURCE & RESEARCH CENTER: UC DAVIS

mmrrc@ucdavis.edu

530-754-MMRRC

Protocol Name: CR11758 Cysrt1 EXDEL

Protocol: GoTaq® G2 Colorless Master Mix(Promega)

Reagent/Constituent	Volume (µL)
Water	4.5
GoTaq® G2 Colorless Master Mix,2X	7.5
Primer 1. (stock concentration is 20µM) comF	0.5
Primer 2. (stock concentration is 20µM) wtR	0.5
Primer 3. (stock concentration is 20µM) mutR	0.5
DNA (example) extracted w/ "Qiagen DNeasy columns or other similar silica based kits"	1.5
TOTAL VOLUME OF REACTION:	15.00 µL

Comments on protocol:

- Protocol may work with other DNA extraction methods.

Strategy:

Steps	Temp (°C)	Time (m:ss)	# of Cycles
1. Initiation/Melting HOT START? <input type="checkbox"/>	94	2:00	1x
2. Denaturation	94	0:10	
3. Annealing steps 2-3-4 cycle in sequence	65 (↓1°C/cycle)	0:30	10x
4. Elongation	68	2:00	
5. Denaturation	94	0:15	
6. Annealing steps 5-6-7 cycle in sequence	55	0:30	25x
7. Elongation	68	2:00 (↑20sec/cycle)	
8. Finish	4	∞	n/a

Primers:

Name	Nucleotide Sequence (5' - 3')	Electrophoresis Protocol:		
1. CR_Cysrt1_comF	GCTGGAGGACTAAGAAGACAGACG	Agarose: 1.5%	V: 90	
2. CR_Cysrt1_wtR*	AGTACCCCTTGATGCCCTTGATACC	Estimated Running Time: 90 min.		
3. CR_Cysrt1_mutR	CCTCTGTAAAACAGCAGTCATGACC	Primer Combination	Band (bp)	Genotype
		1 & 2, 1 & 3	572, 1225	wildtype
		1 & 3	359	mutant

Allele Description: Exon 1 ENSMUSE00001318733 was constitutively deleted from the 5'UTR through the 3'UTR from the Cysrt1 Gene ENSMUSG00000036731 using CRISPR Cas9 gene editing technology in mouse zygotes. The subsequent 866bp deletion from chr2: 25128878- 25129743 GRCm39 was sequence verified. The selected founder animal was backcrossed to C57BL6/N to produce sequence confirmed heterozygous animals for establishing and archiving the line.

*wtR primer untested (ePCR verified)

