

# GENOTYPING PROTOCOL

## MUTANT MOUSE RESOURCE & RESEARCH CENTER: UC DAVIS

[mmrrc@ucdavis.edu](mailto:mmrrc@ucdavis.edu)

530-754-MMRRC

**Protocol Name:** CR11767 Haus8 EXDEL

**Protocol:** GoTaq® G2 Colorless Master Mix(Promega)

| Reagent/Constituent                                                                   | Volume (μL)     |
|---------------------------------------------------------------------------------------|-----------------|
| Water                                                                                 | 4.5             |
| GoTaq® G2 Colorless Master Mix,2X                                                     | 7.5             |
| Primer 1. (stock concentration is 20μM) comF                                          | 0.5             |
| Primer 2. (stock concentration is 20μM) wtR                                           | 0.5             |
| Primer 3. (stock concentration is 20μM) mutR                                          | 0.5             |
| DNA (example) extracted w/ "Qiagen DNeasy columns or other similar silica based kits" | 1.5             |
| <b>TOTAL VOLUME OF REACTION:</b>                                                      | <b>15.00 μL</b> |

### Comments on protocol:

- Protocol may work with other DNA extraction methods.

### Strategy:

| Steps                                                     | Temp (°C )      | Time (m:ss)         | # of Cycles |
|-----------------------------------------------------------|-----------------|---------------------|-------------|
| 1. Initiation/Melting HOT START? <input type="checkbox"/> | 94              | 2:00                | 1x          |
| 2. Denaturation                                           | 94              | 0:10                |             |
| 3. Annealing steps 2-3-4 cycle in sequence                | 65 (↓1°C/cycle) | 0:30                | 10x         |
| 4. Elongation                                             | 68              | 2:00                |             |
| 5. Denaturation                                           | 94              | 0:15                |             |
| 6. Annealing steps 5-6-7 cycle in sequence                | 55              | 0:30                | 25x         |
| 7. Elongation                                             | 68              | 2:00 (↑20sec/cycle) |             |
| 8. Finish                                                 | 4               | ∞                   | n/a         |

### Primers:

| Name             | Nucleotide Sequence (5' - 3') | Agarose: 1.5% V: 90                                        |
|------------------|-------------------------------|------------------------------------------------------------|
| 1. CR_Haus8_comF | GTCATCAGGCTAGGTGACAAGC        | Estimated Running Time: 90 min.                            |
| 2. CR_Haus8_wtR* | GGACTCCACCACTCTTCTTCC         | <b>Primer Combination</b> <b>Band (bp)</b> <b>Genotype</b> |
| 3. CR_Haus8_mutR | CTACCTGGTTTATCAAGGGGATGG      | 1 & 2, 1 & 3 387,689 wildtype                              |
|                  |                               | 1 & 3 395 mutant                                           |

### Electrophoresis Protocol:

**Allele Description:** Exon 3 ENSMUSE00001249928 and flanking splicing regions were constitutively deleted from the Haus8 Gene ENSMUSG00000035439 using CRISPR Cas9 gene editing technology in mouse zygotes. The subsequent 294bp deletion from Chr8:71715628-71715921 GRCm39 was sequence verified. The selected founder animal was backcrossed to C57BL6/N to produce sequence confirmed heterozygous animals for establishing and archiving the line.

\*wtR primer untested (ePCR verified)

